



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Impact of Ebola outbreak on reproductive health services: implementing an ambulance referral system in a rural district of Sierra Leone

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-029093
Article Type:	Research
Date Submitted by the Author:	14-Jan-2019
Complete List of Authors:	<p>Quaglio , GianLuca ; European Parliament  Tognon, Francesca; Department for Woman and Child Health, University of Padua, Italy  Finos, Livio; Department of Developmental Psychology and Socialisation, University of Padua, Italy  Bome, David; Ministry of Health and Sanitation, Freetown, Sierra Leone  Sesay, Santigie ; Ministry of Health and Sanitation, Freetown, Sierra Leone  Atiba, Kebbie ; Department for Woman and Child Health, Pujehun Hospital, Sierra Leone  Di Gennaro, Francesco ; Department of Infectious Diseases, University of Bari, Italy  Bienvenu Salim, Camara ; National Centre for Training and Research in Rural Health of Maferinyah, Forécariah, Guinea  Marotta, Claudia; Department of Science for Health Promotion and Mother to Child Care, University of Palermo, Italy  Pisani, Enzo; Department for Woman and Child Health, Pujehun Hospital, Sierra Leone  Bangura , Zainab; Department for Woman and Child Health, Pujehun Hospital, Sierra Leone  Pizzol, Damiano; Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy  Saracino, Annalisa; Department of Infectious Diseases, University of Bari, Italy  MAZZUCCO, WALTER; Department of Science for Health Promotion and Mother to Child Care, University of Palermo, Italy  Jones, Susan; Department of Nursing and Midwifery School of Human and Health Sciences, University of Huddersfield, UK  Putoto, G; Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy</p>
Keywords:	PAEDIATRICS, Public health < INFECTIOUS DISEASES, Community gynaecology < GYNAECOLOGY

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



# Impact of Ebola outbreak on reproductive health services: implementing an ambulance referral system in a rural district of Sierra Leone

Gianluca Quaglio,<sup>1,2,3</sup> Francesca Tognon,<sup>4</sup> Livio Finos,<sup>5</sup> David Bome,<sup>6</sup> Sesay Santigie,<sup>6</sup> Atiba Kebbie,<sup>7</sup> Francesco Di Gennaro,<sup>8</sup> Bienvenu Salim Camara,<sup>9</sup> Claudia Marotta,<sup>10</sup> Vincenzo Pisani,<sup>7</sup> Bangura Zainab,<sup>7</sup> Damiano Pizzol,<sup>3</sup> Annalisa Saracino,<sup>8</sup> Walter Mazzucco,<sup>10</sup> Susan Jones,<sup>11</sup> Giovanni Putoto<sup>3</sup>

1. Directorate-General for Parliamentary Research Services (EPRS), European Parliament, Brussels, Belgium;
2. Department of International Health/CAPHRI, Maastricht University, The Netherlands;
3. Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy;
4. Department for Woman and Child Health, University of Padua, Italy;
5. Department of Developmental Psychology and Socialisation, University of Padua, Italy;
6. Ministry of Health and Sanitation, Freetown, Sierra Leone;
7. Department for Woman and Child Health, Pujehun Hospital, Sierra Leone;
8. Department of Infectious Diseases, University of Bari, Italy;
9. National Centre for Training and Research in Rural Health of Maferinyah, Forécariah, Guinea;
10. Department of Science for Health Promotion and Mother to Child Care, University of Palermo, Italy;
11. Department of Nursing and Midwifery School of Human and Health Sciences, University of Huddersfield, UK.

## Corresponding author

Gianluca Quaglio, MD, PhD

Directorate-General for Parliamentary Research Services (EPRS)

European Parliament

Rue Wiertz, 60, B-1047

Brussels, Belgium

gianluca.quaglio@europarl.europa.eu

**Words count** (excluding title page, abstract, references, figures and tables): 5182

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**ABSTRACT**

**Objectives** To assess the utilisation of maternal and child health (MCH) services before, during, and after the Ebola virus disease outbreak.

**Design** A prospective observational study of MCH services.

**Setting** Pujehun district in Sierra Leone. Data was collected from 77 community health facilities and 1 hospital over a 6-year period from January 2012 to December 2017.

**Main outcome measures** The utilisation of MCH services was evaluated by assessing: i) institutional deliveries, Cesarean-sections, paediatric and maternity admissions, paediatric and maternity deaths, and major direct obstetric complications, at hospital level; ii) ante natal care 1 and 4, institutional delivery, and family planning, at community level. The contribution of a strengthened referral system was also analysed.

**Results** At hospital level, data between the Ebola period and the pre-Ebola period shows a statistically significant increase in the number institutional deliveries ( $p=0.02$ ) and a reduction of maternal deaths ( $p=0.042$ ). There was statistical significance between the post Ebola vs Ebola period, and post Ebola vs pre-Ebola periods for all indicators considered. At community level, with the exception of family planning, the differences between the Ebola period and pre-Ebola period are statistically significant for all indicators: institutional delivery ( $p < 0.001$ ), ANC 1 ( $p = 0.042$ ), and ANC 4 ( $p = 0.008$ ). The differences between averages of the post Ebola vs pre-Ebola were significant, with an increase for institutional delivery ( $p < 0.001$ ) and ANC 4 ( $p < 0.001$ ). However, there was a statistically significant negative difference between trends in the two periods, for all the variables considered. The RS determined a significant increase in major direct obstetric complications and pediatric cases.

**Conclusions** Due to the strengthened referral service and a stronger health system compared to other districts in Sierra Leone, health facilities in Pujehun district, at community and hospital level, were able to maintain service provision and uptake during and after the Ebola epidemic.

**Keywords:** Ebola, Sierra Leone, Maternal and Child Health indicators, Referral system, Reproductive health service.

## Strengths and limitations of this study

- ▶ The study uses data from a remote rural district in Sierra Leone, with a 6-year observational period. Data have been collected in a prospective way, reducing the potential bias in the accuracy of the data reported by other studies carried out in countries affected by Ebola.
- ▶ The pre, intra, and post-Ebola periods data, allowed a comparison between trends.
- ▶ The data refers to a single area of Sierra Leone: the sample cannot be considered representative of the country as a whole.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**INTRODUCTION**

The 2014-2015 Ebola Virus Disease (EVD) outbreak was the most severe in history, mainly affecting three West African countries; Guinea, Sierra Leone and Liberia. Overall 28,616 people were infected of which 11,310 died and the outbreak was declared a global public health emergency by the WHO.<sup>1</sup> Of the three countries affected, Sierra Leone had the most confirmed cases (8,704), which accounted for 50% of all confirmed cases in West Africa, and 3,589 deaths.<sup>2-4</sup> All 14 districts in Sierra Leone were affected, but at different times and to varying degrees.<sup>5</sup> During the Ebola crisis the population’s trust in the national health system declined in Sierra Leone, leading to an overall reduction in the use of health services, including reproductive, maternal, and child services.<sup>6-8</sup> Underlying factors for the decrease in the use of health services included fear of infection, for both healthcare workers and patients, the underlying fragility of the health systems, the reduced numbers of available health personnel, and the death of healthcare workers due to EVD.<sup>9-10</sup> It has been estimated that 30% of health workers who died of EVD in West Africa were maternal and child healthcare (MCH) providers.<sup>11</sup> However, there were considerable variations in the reduction of health service uptake when looked at by district level in Sierra Leone.<sup>6-12-14</sup> While districts such as Kambia, Port Loko and Bonthe showed large reductions in facility-based delivery (between 38-41%), the district of Pujehun showed only a 5% decrease in the same service. Similar geographic variations were seen in the reduction in antenatal care (ANC) visits.<sup>12-13</sup>

The number of confirmed EVD cases - and deaths - varied considerably by district. There were no more than 100 confirmed cases in both Bonthe and Pujehun, and up to 4,000 confirmed cases in both Port Loko and Bombali.<sup>15</sup> However, public fear of Ebola, regardless of the actual number of cases per district, may still have prevented many people from accessing services. The challenge of providing adequate levels of care during a humanitarian emergency such as the EVD crisis was further exacerbated by the weak health system in Sierra Leone, particularly in rural areas where the poor condition of the roads and high transport costs cause delays in accessing services, and contribute to increased maternal and neonatal mortality.<sup>16</sup> Different types of referral systems (RSs) such as motorbikes were present in the country in the pre Ebola period to transport patients from the villages to the nearest health facility. Ambulances were also present in several districts with 73% of health facilities nationwide having a functioning RS, 59% of them consisting of an ambulance on call.<sup>12-17</sup> In the Pujehun district, the RS was barely functioning, only able to support the activity of a limited number of Peripheral Health Units (PHUs). The service was also entirely

1  
2  
3 funded by the patients themselves, resulting in underutilization of the service. Utilization was  
4 further reduced during the outbreak, when the ambulances were identified by the population with  
5 the transport of Ebola infected patients, and their use occasioned fear and distrust.  
6  
7

8 Doctors with Africa (DwA) CUAMM is an Italian NGO working in Sierra Leone since 2012. It  
9 is present in the Pujehun district focusing on MCH care both at hospital and community level.<sup>18</sup> In  
10 January 2015, in collaboration with the Ministry of Health and Sanitation (MoHS) of Sierra Leone  
11 and UNICEF, DwA began the re-organisation and reinforcement of the RS, transferring pregnant  
12 women and pediatric cases from PHUs to the Pujehun hospital. Our previous study carried out in  
13 the district of Pujehun showed that activities undertaken to manage the EVD outbreak and  
14 preserve MCH services at the district hospital and at the community level reduced the spread of  
15 infection and the impact of the disease on MCH services.<sup>18</sup> As widely reported in our previous  
16 study, the approach implemented in the Pujehun district avoided vertical interventions: it worked  
17 on strengthening all the components of the health system - governance, human resources,  
18 community involvement - before, during, and after the epidemic. The previous study<sup>18</sup> provided  
19 information only on three MCH indicators, namely pediatric admissions, maternity admissions, and  
20 institutional deliveries; in addition it did not assess the trends in the post-EVD period. Existing  
21 studies examining the influence of EVD on MCH services targeted the outbreak and the immediate  
22 post-outbreak periods.<sup>19-22</sup> Understanding the trends in the use of MCH services before, during,  
23 and after the EVD outbreak will help to guide post-EVD interventions, increasing access to MCH  
24 services in rural Sierra Leone. This information will also be useful in preparing a more organised  
25 and structured RS. With this background, the aims of this study are: i) to assess institutional  
26 deliveries, C-sections, paediatric and maternity admissions, paediatric and maternity deaths, and  
27 major direct obstetric complications (MDOCs), before, during, and after the EVD in the Pujehun  
28 hospital, thus complementing the results of the previous report which were limited to 3 MCH  
29 indicators; ii) to assess the use of ANC 1 and 4, institutional delivery, and family planning, at  
30 community level. This study was carried out in conjunction with the strengthening of an RS  
31 initiated a few weeks after the Pujehun district was declared Ebola-free.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**METHODS**

**Setting**

Sierra Leone has four provinces that are divided into 14 districts. Pujehun is one of four districts in the southern province. It has a population of approximately 375,000 inhabitants. The primary care network included 77 MoHS PHUs, 5 of which provide basic emergency obstetrics care (BEmOC). The secondary care system consists of the MoHS provided district hospital, which comprises the MCH complex, providing comprehensive emergency obstetric and newborn care (CEmONc) services. Connections between the community and health facilities are difficult because of the very poor condition of the roads. Furthermore, the district is divided by a major river (Moa River) and has a riverine area reachable only with boats, which further hinders access. The first case of Ebola in Pujehun district was reported on the 7th July 2014. The district was declared Ebola free on the 10th January 2015.<sup>23</sup> A total of 49 patients were registered with a case fatality rate of 85.7% (42/49).

**Referral system**

In the Pujehun district, two ambulances managed by the District Health Management Team (DHMT) were functioning in the pre Ebola period, but only 63% of the PHUs were able to use the service.<sup>12 17</sup> Emergency calls were not coordinated by the hospital and the transport costs were covered by the patients, dissuading many from using the service. During the outbreak, people came to associate the ambulances with transporting Ebola infected patients, which further discouraged their use. A 24-h free-of-charge ambulance RS, transferring pregnant women with obstetric complications from the health centers to Pujehun hospital was implemented in January 2015. In the hospital a call center was established and the call center number was distributed to all the 77 PHUs. Private calls were considered only in the case of an emergency or if the staff of the PHU were not available. After confirming an emergency condition together with the PHU staff, the hospital midwife had the responsibility to authorize the referral. A nurse on duty from the maternity hospital accompanied the driver in each referral. PHU staff were trained together with the hospital staff and DHMT to recognize and manage obstetric emergencies. All healthcare workers involved in the emergency transfer system received regular feedback on the appropriateness of each referral carried out. Referrals were carried out by 3 ambulances, two positioned in the Pujehun MCH complex, and a third one in Jendema, bordering Liberia, on the

opposite side of the Moa River. Around the Jendema area, 15 PHUs were located serving a population of approximately 80,000 inhabitants. Referrals in this area were made using the ambulances and by transferring patients at the river crossing point via a barge or a motor boat, depending on the flow rate of the river. Pediatric referrals were performed using private motor bikes available in the villages and hired from PHUs staff without the involvement of the call center. A referral form describing the clinical case and the justification for the referral was distributed to all the PHUs. The bike rider, after bringing the patient to the pediatric ward, delivered the referral form and received the reimbursement. For all patients carried to the hospital information was collected, including demographics, location, and the reason for contacting the RS. Community awareness activities were organized about the RS through meetings and radio discussions held by the DMHT, hospital health personnel, and local authorities.

### **Study design, population, and period**

A prospective observational study using routinely collected health services data, from January 2012 to December 2017, was carried out. Three time periods were considered: pre- Ebola period (1<sup>st</sup> January 2012 – 30<sup>th</sup> May 2014); Ebola period (1<sup>st</sup> June 2014 – 28<sup>th</sup> February 2015); post- Ebola period (1<sup>st</sup> March 2015 – 31<sup>th</sup> December 2017). We considered the Ebola period from one month before the first confirmed case in the district (i.e. June 2014), to three months after the last confirmed case in the district (i.e. February 2015). This was done because in Sierra Leone the outbreak had started in other districts of the country before the first case registered in Pujehun and continued to affect other districts until November 2015. It is realistic to assume that public fear of potential EVD cases and lack of confidence in the health services persisted in the Pujehun population during that time.<sup>14</sup> In addition, expanding the Ebola period enabled a full assessment of the impact of the disease with an adequate comparison with the two long periods before and after the Ebola epidemic.

### **Data collection**

Data on MCH indicators was prospectively collected from hospital registers (maternity ward, delivery unit, pediatric ward, operating theatre) directly by DwA. The following variables were collected: 1) paediatrics admissions per month; 2) pediatric deaths per month; 3) maternity admissions per month; 4) maternal deaths per month; 5) deliveries per month; 6) C-sections per

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

month; 7) MDOC cases per month. At community level, the following variables were collected from the local district Health Management Information System, with the technical support of DWA: 1) family planning consultations per month; 2) deliveries per month; 3) ANC 1 per month; 4) ANC 4 per month. Different variables were collected from the two types of sites, based on the different services provided at community level (BEmONC) and at hospital level (CEmONC). For the RS, data was collected from records of all of the study sites, including delivery registers, delivery logbooks, prenatal registers, referral registers, and death registers. Additional data was collected from the ambulance database and logbook. Records in the database were then validated by cross-checking the records with registers at the study sites.

**Statistical analysis**

For each indicator, a segmented seasonal autoregressive model of order 1 was estimated. The segments defined the three periods: before the EVD epidemic (January 2012 to May 2014), during the epidemic (June 2014 to February 2015), and after the epidemic (March 2015 to December 2017). The model for each indicator  $Y_t$  collected at hospital or community level was as follows:  $Y_t = \beta_0 + \beta_1 T_t + \beta_2 X_t + \beta_3 X_t T_t + \beta_4 Z_t + \beta_5 Z_t T_t + \beta_6 Month + \varepsilon_t$ .  $\beta_0$  estimates the number of individuals using the service at the beginning of the pre-Ebola period;  $\beta_1$  estimates the average monthly change in the number using the service over the pre-outbreak period;  $T_t$  is the time since the start of the study;  $\beta_2$  represents the change in the level of service use that occurred in the period immediately after the EVD period (designated by indicator variable  $X_t$ );  $\beta_3$  represents the difference between the trend in service use during the EVD outbreak compared to the pre-disease period;  $\beta_4$  represents the change in service use that occurred in the period immediately after the end of the outbreak (post-outbreak period designated by indicator variable  $Z_t$ );  $\beta_5$  is the difference between the trend in service use during the period after the Ebola virus disease outbreak compared with the period during the outbreak period;  $\beta_m$  represents a series of indicator variables for each calendar month, and  $t$  is the random error term. Overall trends across the periods and the comparisons among trends were calculated as follows: linear trend during the outbreak =  $\beta_1 + \beta_3$ ; linear trend after the outbreak =  $\beta_1 + \beta_3 + \beta_5$ ; and linear trend after the outbreak vs linear trend before the outbreak =  $\beta_3 + \beta_5$ . Average levels across the periods and their comparisons were calculated as follows: average during the outbreak =  $\beta_0 + \beta_2$ ; average after the outbreak =  $\beta_0 + \beta_2 + \beta_4$ ; and difference between after the outbreak and before the

outbreak =  $\beta_2 + \beta_4$ . Differences were considered statistically significant at  $p < 0.05$ . The analysis was performed using R.<sup>24</sup>

### Patient involvement

No patients were involved in defining the research question or the outcome measures, nor were they involved in the design and implementation of the study. There are no plans to involve patients in the dissemination of the results.

## RESULTS

### Hospital level: Pre-Ebola period

At hospital level, the pre Ebola period for MCH indicators showed an average of 49 maternal admissions per month (95% CI 37 to 61,  $p < 0.001$ ), 9 C-sections per month (95% CI 4 to 14,  $p = 0.001$ ), and 16 MDOCs per month (95% CI 5 to 26,  $p = 0.003$ ). There were an average of 46 pediatric admissions per month (95% CI 10 to 82,  $p = 0.011$ ), and 27 institutional deliveries per month (95% CI 19 to 34,  $p < 0.001$ ). For all indicators, the trend is stable during the pre Ebola period, without significant changes (Table 1 and 2; Figure 1 and 2).

### Hospital level: Ebola vs pre-Ebola period

At hospital level, the differences between Ebola period vs pre-Ebola averages show a statistically significant increase for institutional deliveries (11, 95% CI 2 to 21,  $p = 0.02$ ) and for the reduction of maternal deaths (-1, 95% CI - 2 to 0,  $p = 0.042$ ). There is also a statistically significant difference between the trend of Ebola period vs pre-Ebola period, for maternal admissions (7, 95% CI 4 to 11,  $p < 0.001$ ), MDOCs (4, 95% CI 1 to 7,  $p = 0.006$ ), and institutional deliveries (4, 95% CI 2 to 6,  $p = 0.001$ ) (Table 1 and 2; Figure 1 and 2).

### Hospital level: Ebola vs post-Ebola period

At hospital level, the differences between averages of the post Ebola vs Ebola are statistically significant for all indicators: institutional deliveries, C-sections, paediatric and maternity admissions, paediatric and maternity deaths, and MDOCs. There is also a negative trend in the transition from Ebola to post Ebola for maternal admissions (-7, 95% CI -10 to -4,  $p < 0.001$ ),

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

MDOCs (-4, 95% CI -7 to -1, p 0.009) and institutional deliveries (-3, 95% CI -5 to -1, p 0.001) (Table 1 and 2; Figure 1 and 2).

**Hospital level: Post-Ebola vs pre-Ebola period**

The differences between averages of the post Ebola vs pre-Ebola periods are also statistically significant for all indicators, except for maternal deaths. The differences between trends between post-Ebola vs pre-Ebola period are only significant for pediatric admissions (3, 95% CI 0 to 5, p 0.035) (Table 1 and 2; Figure 1 and 2).

**Table 1** Maternal admissions, maternal deaths, C-sections, and MDOCs at hospital level

	Maternal admissions			Maternal deaths			C-sections			MDOC		
	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value
Difference between average of Ebola period vs pre-Ebola period	7	-7 to 22	0.333	-1	-2 to 0	0.042	5	-1 to 11	0.13	2	-11 to 14	0.782
Difference between average of post-Ebola period vs Ebola period	43	28 to 58	<0.001	2	1 to 3	0.001	15	8 to 21	<0.001	41	30 to 54	<0.001
Difference between average of post-Ebola period vs pre-Ebola period	50	37 to 64	<0.001	1	0 to 2	0.135	19	13 to 25	<0.001	43	31 to 54	<0.001
<b>Pre-Ebola period</b>												
Number of events over pre-Ebola period ( $\beta_0$ )	49	37 to 61	<0.001	1	0 to 2	0.026	9	4 to 14	0.001	16	5 to 26	0.003
Trend in number over pre-Ebola period ( $\beta_1$ )	0	0 to 1	0.281	0	0 to 0	0.677	0	-0 to 0	0.999	0	0 to 0.5	0.768
<b>Ebola period</b>												
Average monthly change in number over Ebola period ( $\beta_2$ )	-40	-60 to -19	<0.001	0	-2 to 0	0.480	2	-7 to 11	0.668	-11	-29 to 6	0.207
Difference between trend of Ebola period vs pre-Ebola period ( $\beta_3$ )	7	4 to 11	<0.001	0	0 to 0	0.605	1	-1 to 2	0.346	4	1 to 7	0.006
<b>Post-Ebola period</b>												
Average monthly change in number during post-Ebola period ( $\beta_4$ )	11	-7 to 30	0.23	1	0 to 2	0.258	13	5 to 21	0.001	16	0 to 32	0.044
Difference between trend of post-Ebola period vs Ebola period ( $\beta_5$ )	-7	-10 to -4	<0.001	0	0 to 0	0.665	-1	-2 to 0.8	0.433	-4	-7 to -1	0.009
Difference between trend of post-Ebola vs pre-Ebola period ( $\beta_3 + \beta_5$ )	0	-1 to 1	1	0	0 to 0	0.657	0	0 to 0	0.431	0	0 to 1	0.503

**Table 2** Pediatric admissions, pediatric deaths, and institutional deliveries at hospital level

	Pediatric admissions			Pediatric deaths			Institutional deliveries		
	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value
Difference between average of Ebola period vs pre-Ebola period	1	-39 to 40	0.968	-1	-6 to 5	0.826	11	2 to 21	0.02
Difference between average of post-Ebola period vs Ebola period	133	92 to 174	<0.001	9	3 to 15	0.004	28	18 to 38	<0.001
Difference between average of post-Ebola period vs pre-Ebola period	134	98 - 170	<0.001	8	3 to 14	0.003	39	31 to 48	<0.001
<b>Pre-Ebola period</b>									
Number of events over pre-Ebola period ( $\beta_0$ )	46	10 to 82	0.011	7	2 to 12	0.007	27	19 to 34	<0.001
Trend in number over pre-Ebola period ( $\beta_1$ )	0	-2 to 2	0.808	0	0 to 0	0.641	0	0 to 0	0.42
<b>Ebola period</b>									
Average monthly change in number over Ebola period ( $\beta_2$ )	1	-48 to 50	0.955	1	-7 to 9	0.836	-12	-25 to 1	0.072
Difference between trend of Ebola period vs pre-Ebola period ( $\beta_3$ )	1	-8 to 10	0.823	0	-1 to 2	0.763	4	2 to 6	0.001
<b>Post Ebola period</b>									
Average monthly change in number over post-Ebola period ( $\beta_4$ )	53	5 to 100	0.029	6	-1 to 14	0.086	11	-1 to 22	0.064
Difference between trend of post-Ebola period vs Ebola period ( $\beta_5$ )	2	-7 to 10	0.702	0	-1 to 1	0.899	-3	-5 to -1	0.001
Difference between trend of post-Ebola vs pre-Ebola period ( $\beta_3 + \beta_5$ )	3	0 to 5	0.035	0	0 to 0	0.423	0	0 to 0	0.486

Table 3 Institutional delivery, ANC 1, ANC 4 and family planning at community level

	Institutional delivery			ANC 1			ANC 4			Family planning		
	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value
Difference between average of Ebola period vs pre-Ebola period	148	99 to 196	<0.001	74	3 to 145	0.042	80	21 to 139	0.008	490	-92 to 1073	0.099
Difference between average of post-Ebola period vs Ebola period	-10	-59 to 39	0.695	-48	-122 to 26	0.2	23	-38 to 84	0.461	-262	-855 to 330	0.386
Difference between average of post-Ebola period vs pre-Ebola period	138	93 to 183	<0.001	26	-40 to 91	0.448	103	48 to 157	<0.001	228	-293 to 750	0.391
<b>Pre Ebola period</b>												
Number of events over pre-Ebola period ( $\beta_0$ )	688	643 to 732	<0.001	1062	1002 to 1121	<0.001	694	644 to 743	<0.001	2690	2187 to 3193	<0.001
Trend in number over pre-Ebola period ( $\beta_1$ )	8	6 to 10	<0.001	7	4 to 10	<0.001	6	4 to 8	<0.001	69	42 to 95	<0.001
<b>Ebola period</b>												
Average monthly change in number over Ebola period ( $\beta_2$ )	-28	-90 to 34	0.382	-61	-161 to 40	0.238	-94	-176 to -11	0.027	-671	-1431 to 89	0.084
Difference between trend of Ebola period vs pre-Ebola period ( $\beta_3$ )	-1	-12 to 10	0.881	-5	-21 to 12	0.591	5	-8 to 19	0.437	-26	-156 to 104	0.692
<b>Post Ebola period</b>												
Average monthly change in number during post-Ebola period ( $\beta_4$ )	-25	-81 to 30	0.37	-5	-94 to 83	0.906	35	-37 to 109	0.343	-51	-759 to 657	0.888
Difference between trend of post-Ebola period vs Ebola period ( $\beta_5$ )	-7	-17 to 4	0.228	-2	-18 to 15	0.819	-13	-27 to 0	0.056	-59	-186 to 68	0.361
Difference between trend of post-Ebola vs pre-Ebola period ( $\beta_3 + \beta_5$ )	-7	-10 to -4	<0.001	-6	-10 to -3	<0.001	-8	-11 to -5	<0.001	-85	-119 to -51	<0.001

### Community level: Pre-Ebola period

At community level, all the maternal health indicators in the months before Ebola showed a positive trend. There was a monthly average increase of 8 institutional deliveries (95% CI 6 to 10,  $p < 0.001$ ); a monthly average increase of 7 ANC 1 (95% CI 4 to 10,  $p < 0.001$ ) and 6 ANC 4 (95% CI 4 to 8,  $p < 0.001$ ), and a monthly average increase of 69 women accessing family planning services (95% CI 42 to 95,  $p < 0.001$ ) (Table 3; Figure 3).

### Community level: Ebola vs pre-Ebola period

At community level, with the exception of family planning, the differences between averages of Ebola period vs pre-Ebola are statistically significant for all indicators: institutional deliveries (148, 95% CI 99 to 196,  $p < 0.001$ ), ANC 1 (74, 95% CI 3 to 145,  $p = 0.042$ ), and ANC 4 (80, 95% CI 21 to 139,  $p = 0.008$ ). The average monthly change in number during the Ebola period was negative for the 4 indicators considered, but statistically significant only for the ANC 4 (-94, 95% -176 to -11,  $p = 0.027$ ). There was no statistically significant difference between the Ebola period and pre-Ebola trends for any of the indicators (Table 3; Figure 3).

### Community level: Ebola vs post-Ebola period

At community level, the differences between averages and the difference between trends of the post Ebola vs Ebola period are not significant for any of the indicators considered (Table 3; Figure 3).

### Community level: Post Ebola vs pre-Ebola period

The differences between averages of the post Ebola vs pre-Ebola are statistically significant, with an increase in institutional deliveries (138, 95% CI 93 to 183,  $p < 0.001$ ) and ANC 4 (103, 95% CI 48 to 157,  $p < 0.001$ ) (Table 3; Figure 3). However, there is a negative difference between trends among the two periods, for all the variables considered: institutional deliveries (-7, 95% CI -10 to -4,  $p < 0.001$ ) ANC 1 (-6, 95% CI -10 to -3,  $p < 0.001$ ), ANC 4 (-8, 95% CI -11 to -5,  $p < 0.001$ ) and family planning (-85, 95% CI -119 to -51,  $p < 0.001$ ) (Table 3; Figure 3).



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Referral system: Obstetric and paediatric results**

Between January 2015 and December 2017 there were 2,450 obstetric referrals. Of these, 1,574 (64%) were MDOC, which represent 70% of all the 2,233 MDOCs treated in the hospital over the same period. The baseline characteristics and reasons for MDOCs collected through the RS are reported on Table 4. At the same time, 4,671 paediatric patients were admitted in the hospital through the RS, representing 72% of the 6,518 total admission during the same period. Reasons for paediatric referrals are shown on Table 5.

**Table 4** Baseline characteristics and reasons for MDOCs collected through RS, period 2015 - 2017

Age (years)	N	%
Mean	25,3	SD 7
12-19	442	28%
20-29	613	39%
30-39	464	29%
40+	43	3%
Unknown	12	1%
Number of previous deliveries		
0	474	30%
1 or 2	377	24%
3 or 4	292	19%
5 or 6	207	13%
7+	212	13%
Unknown	12	1%
MDOC treated		
Prolonged/obstructive labour	848	54%
Antepartum haemorrhage	195	12%
Severe pre-eclampsia/eclampsia	165	11%
Abortum complicatum	117	7%
Post-partum haemorrhage	157	10%
Ectopic pregnancy	24	2%
Rupture uterus	30	2%
Sepsis	38	2%
Total	1574	100%

**Table 5** Reasons for paediatric RS, period 2015-2017\*

Reason for referral	Number	%
Malaria	1540	30%
Anemia	910	18%
Pneumonia/ARI**	830	16%
Diarrhoea and vomiting	495	10%
Malnutrition	274	5%
Convulsion	186	4%
Hernia/Hydrocele	165	3%
Sepsis/Septicemia	127	2%
Dehydration	48	1%
Burn	30	1%
Others	522	10%
Total	5127	100%

\* For a number of patients, more than one suspected diagnosis for referral was reported; \*\* Acute Respiratory Infection

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**DISCUSSION**

This study presents for the first time trends in utilization of MCH services before, during, and after Ebola, at hospital and community level from the country most heavily affected by the Ebola epidemic. It also presents data on the restructured and reorganised RS, which started immediately after the EVD outbreak. The study shows that there was a decrease in all MCH indicators and service uptake immediately after the onset of the outbreak, with a levelling or increase during the EVD period. In the post-Ebola period, all indicators (except for maternal deaths) showed an increase, in comparison with the pre-Ebola period. This was particularly marked at hospital level because the post Ebola reinforcement of the RS led to an increase in pediatric admissions, maternal admissions, and consequently a rise of institutional deliveries, C-sections, and MDOCs. In addition, while at the hospital level trends in the post-Ebola period are in line with the pre-Ebola, at community level there is a negative trend compared to the pre-Ebola period for all indicators taken into consideration. The study presents results in contrast to other studies that showed a decline in MCH services at facility and community levels in the Ebola and post-Ebola periods.<sup>6 25 26</sup>

**Pre Ebola and Ebola periods**

As extensively described in our previous reports,<sup>18 27</sup> a number of measures were put in place to control the Ebola epidemic in the Pujehun district which reduced the impact of the disease on mothers and children compared to other districts. During the EVD epidemic, the focus on vertical programmes was frequently associated with failures in basic management measures for controlling a disease outbreak. Rather than vertical interventions, the approach implemented in the Pujehun district focussed on all the components of the health system, beginning before the EVD crisis. A rapid response to the crisis by the local health authorities was implemented adopting public health measures before any other district in Sierra Leone.<sup>28</sup> The activities were mainly concentrated on keeping the health service open and properly functioning in order to reduce the collateral effects of the epidemic on routine health services. No health units in the Pujehun district were closed during the epidemic. Measures to empower community leaders and use culturally appropriate methods of communication helped to dispel community mistrust in the health services. At community level, a number of strategies were implemented such as the regular rotation of health facility staff, which strengthened teamwork and effective leadership. In Sierra Leone, healthcare workers based at community health centres may often work alone in isolated

centres with limited support from clinical colleagues or management. By rotating staff through the various facilities, they gain on the job training, peer support, and develop new working relationships. At the start of the Ebola epidemic, many expatriate healthcare workers in NGOs left Sierra Leone, negatively affecting care delivery and staff morale. The continued presence of international teams in the daily activities in Pujehun hospital and the acceptance of the professional risks by both national and international staff may have contributed to maintaining an attitude of 'normality' in an extremely stressful environment. This might also help to explain the population's positive receptiveness towards the health services.<sup>18 27</sup>

At community level, this report showed that family planning, ANC, and institutional deliveries, were affected only at the beginning of the Ebola outbreak with a small decrease in service utilization. In contrast, Jones et al., evaluated the number of antenatal and postnatal visits, institutional births, emergency obstetric care (EmOC), maternal deaths and stillbirths across 13 districts of Sierra Leone for 10 months during, and 12 months prior to the epidemic. They found that following the onset of the epidemic there was an 18% decrease in the number of women attending ANC visits and an 11% decrease in the number of women attending for birth at healthcare facilities.<sup>14</sup>

During the Ebola epidemic, the Pujehun hospital maintained C-sections and delivery volume at pre-Ebola levels. There was a stable number of patients attending the hospital during the Ebola outbreak, as shown by the number of maternal and pediatric admissions. The study of Brolin and colleagues focused on in-hospital deliveries and C-section volume in Sierra Leone. They showed that nationwide, albeit with substantial variation between districts, in-hospital deliveries and C-sections decreased by over 20% during the Ebola outbreak, mainly because of the closure of not-for-profit hospitals.<sup>6</sup> Brolin also noted that in general, at hospital level, in Sierra Leone those facilities that remained open performed about the same number of deliveries and C-sections after the onset of the EVD outbreak as they did before.<sup>6</sup> This seems to indicate that the decrease observed at national level was related to the closing of key health facilities. The number of Ebola cases was not uniform throughout districts in Sierra Leone and Pujehun was one of the least affected districts. The low number of cases may also have helped to maintain public confidence in service provision and uptake of services.<sup>7 8</sup>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Post Ebola period**

There is a shortage of data in Sierra Leone and the other West Africa countries affected regarding the resumption of services after the epidemic. Pujehun district showed contrasting results at community level. Results of the post Ebola vs pre-Ebola show an increase of activities for institutional delivery and ANC 4. However, there is a negative trend among the two periods, for the variables taken into consideration, namely institutional deliveries (-7, 95% CI -10 to -4,  $p < 0.001$ ) ANC 1 (-6, 95% CI -10 to -3,  $p < 0.001$ ), ANC 4 (-8, 95% CI -11 to -5,  $p < 0.001$ ) and family planning (-85, 95% CI -119 to -51,  $p < 0.001$ ). In 2017, the Pujehun district showed a coverage of 98% for ANC 1 (98% in 2013), 91% for ANC 4 (76% at national level in 2013) and 90% for institutional deliveries (62% in 2013).<sup>29 30</sup> The initial intervention carried out by DWA in the period 2012-2014 at the community level probably increased these percentages, with an initial growth of the trend that had been slowing down in the years 2016-2017.

A study by Camara et al. in a rural district of Guinea showed a considerable recovery gap in the post-Ebola period for ANC (37%) and institutional deliveries (34%).<sup>25</sup> Also Delamou et al. noted a significant reduction in the average number of ANC visits and institutional deliveries during the Ebola outbreak, in 6 districts of Guinea, and the overall post-outbreak trends did not suggest recovery.<sup>26</sup> By contrast, Wagenaar et al., which analysed 10 primary care indicators in Liberia, before, during, and after the Ebola outbreak, showed significant positive trends during the post-EVD period for ANC and institutional deliveries.<sup>31</sup> There are multifactorial and complex reasons for the decline of family planning in the Pujehun district. The activities that MoHS and DWA implemented from 2012 onwards were maintained during and after the EVD epidemic. However, a general decrease in the availability of healthcare personnel and international aid was observed. A possible stock-out of family planning methods has also been suggested as a reason for the decrease.<sup>25</sup> In addition, a reduction in demand for family planning in the post Ebola period could account for the decline of the service. Experiencing a disaster can trigger the desire to “rebuild” communities, reducing the need for family planning methods,<sup>32</sup> or communities may prefer traditional methods of contraception.<sup>33</sup> However, the reduction in family planning use in Pujehun district did not translate into an increase in institutional deliveries as occurred in neighbouring Liberia.<sup>34</sup> Although no further transmissions of Ebola took place in the Pujehun district after November 2015, the awareness of the ongoing transmission elsewhere in Sierra Leone, in Guinea and Liberia might have influenced health seeking behaviours.<sup>35 36</sup> However, this does not seem to have influenced other types of MCH services at community level. For comparison, the above

mentioned study of Camara et al. showed that the utilization of family planning declined by 51% during the Ebola outbreak but recovered in the post-Ebola period.<sup>25</sup>

At hospital level, the situation is different. In the post-Ebola period, there was a significant increase in the volumes of activities: pediatric and maternal admissions, MDOC cases, deliveries, and C-sections. This increase can be directly linked to the reorganization and strengthening of the RS immediately after the Ebola epidemic. Based on the 3 delays theory,<sup>37</sup> in Pujehun it was decided to tackle the second delay, a lack of accessibility to health services. The distance to the hospital as well as lack of accessible and affordable vehicles were recognized as significant barriers when attempting to access CEmONC services at the hospital.<sup>38 39</sup> The success of the RS service can be linked to the integration of the key components needed for a successful service, namely: i) a transport system which took account of the specific geographical characteristics of the district;<sup>38</sup> ii) an effective communication system with a call center in contact with all PHUs of the district, the ambulance drivers, and the hospital; iii) training of all the PHU staff on the recognition of obstetric emergencies and on the RS.<sup>40 41</sup> Several meetings were planned with local community leaders and religious leaders to raise awareness of the importance of giving birth in health facilities. Prohibitive costs have been shown to be a major factor in preventing women accessing health facilities during childbirth in Sierra Leone.<sup>38 42 43</sup> Meetings were also organised to inform the population that the service was free of charge, and to give reassurance that the ambulances carried no risk of Ebola infection to people using them. The increase in complicated cases treated at the hospital did not translate into an increase in maternal and pediatric deaths, reflecting positively on the quality of care provided. The maternity ward death rate remained around 1% throughout the 2012-2017 study period. The differences in average death rates during the period 2015-2017 among referred and not referred pediatric patients were 10.5% and 4.3% respectively. This showed that the pediatric RS works for the most critical cases able to reach the hospital in time.

## CONCLUSIONS

There are a number of limitations to this study. The data refers to a single area of Sierra Leone and therefore our sample cannot be considered representative of the country as a whole. We defined our distinct period of EVD outbreak arbitrarily, from one month before the first case in the district to three months after the last case in the district. This was done because the EVD crisis affected areas of the country outside Pujehun prior to and after outbreak within Pujehun. The official end of the EVD epidemic for Sierra Leone was declared on March 17, 2016, and for the countries of

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Guinea and Liberia was declared on June 1, 2016. Finally, our study assumed that no other interventions in addition to those described occurred concurrently with the Ebola epidemic.<sup>18</sup> Similarly, we assumed that no other substantial interventions in addition to the re-organisation of the RS happened in the post-Ebola period which would have affected the service trends that we observed. The strength of this study is that it uses data from a remote rural district in Sierra Leone, with a 6-year observational period. The pre, intra, and post-Ebola periods data, allowed a comparison between trends. DWA was working in this community before the outbreak began, which gave an advantage of knowledge of the setting when the epidemic began, which in turn facilitated mitigating measures to be put in place. In addition, this allowed a collection of data in a prospective way, reducing the potential bias in the accuracy of the data reported by other studies.<sup>6 14 26 31</sup>

During the EVD epidemic, the focus on vertical programmes was frequently associated with failures in basic management procedures for controlling a disease outbreak.<sup>44</sup> The approach implemented in the Pujehun district was not based on vertical interventions: on the contrary it worked on strengthening all the components of the health system - governance, human resources, community involvement - before, during and, after the epidemic. The strengthening of the health system in the district, compared to other districts, allowed the containment of the epidemic and, above all, to maintain and strengthen MCH services as shown by the data reported in the paper. Health facilities in the district, both at community and hospital level, were able to maintain their services during the epidemic, overcoming public fear of Ebola and lack of confidence in service providers, which led to the public staying away from facilities in other districts in Sierra Leone.<sup>14</sup> In post-crisis situations, "windows of opportunity" are opened for redirecting the policies of the national health systems, renovating specific sectors (e.g. human resources, epidemiological surveillance systems, financing, etc.) and renewing services/practices at the operational level.<sup>45</sup> In Pujehun the implementation of an RS immediately after the acute Ebola phase reduced delays in patients accessing care and enabled a significant improvement in all MCH indicators at hospital level. Other studies have also found that using this window of opportunity to introduce systems such as performance based financing can also produce positive outcomes.<sup>46</sup> As Sierra Leone continues its recovery, there is a need to quantify the impact of the outbreak on MCH care to guide long-term strategies for MHC services. This study provides evidence on strategies to increase the resilience of fragile healthcare services and the importance of NGOs and government collaboration to bring about change.

**Acknowledgements** We are deeply grateful to all the staff of Pujehun district hospital and Peripheral Health Units, District Health Management, and the personnel of Doctors with Africa CUAMM who worked in Pujehun during the epidemic. We thank James Dean for his help with proof-reading.

**Contributors** GLQ, FT, and GP contributed to study design, literature review, data analysis, data interpretation, writing, and review of the final manuscript. LF, FDG, DP, and CM performed data analysis, data interpretation, and drafting of the manuscript. DB, SS, AK, BZ, and VP contributed to data collection and data interpretation. BSC, AS, WM, and SJ contributed to data interpretation and review of the final manuscript. All coauthors contributed to the improvement of the article.

**Funding** This work was undertaken and funded by Doctors with Africa CUAMM.

**Competing Interests** None declared.

**Patient consent for publication** Obtained.

**Ethics approval** Sierra Leone Ethics and Scientific Review Committee, Directorate of Policy, Planning and Information, Ministry of Health and Sanitation, Sierra Leone.

**Data sharing statement** All data underlying the findings described in the manuscript are fully available without restriction.

**Disclaimer** The views expressed in this publication are the sole responsibility of the authors and do not necessarily reflect the views of the affiliated organisations.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

REFERENCES

1. World Health Organization. *Ebola response roadmap. Situation reports*. WHO, 2018. Available from: <http://www.who.int/csr/disease/ebola/situation-reports/archive/en/> [Accessed Dec 2018].

2. Jones S, Ameh C. Exploring the impact of the Ebola outbreak on routine maternal health services in Sierra Leone, 2015. Available from: [https://www.vsointernational.org/sites/vso\\_international/files/vso\\_sierra\\_leone\\_\\_](https://www.vsointernational.org/sites/vso_international/files/vso_sierra_leone__) [Accessed Dec 2018].

3. Streifel C. *How did Ebola impact maternal and child health in Liberia and Sierra Leone? A report of the CSIS Global Health Policy Center*. CSIS, 2015.

4. Brolin Ribacke KJ, Saulnier DD, Eriksson A, *et al*. Effects of the West Africa Ebola Virus Disease on Health-Care Utilization - A Systematic Review. *Front Public Health* 2016;4:222.

5. Government of Sierra Leone, Ministry of Health and Sanitation. *Ebola virus disease situation report*. MoHS, 2015.

6. Brolin Ribacke KJ, van Duinen AJ, Nordenstedt H, *et al*. The Impact of the West Africa Ebola Outbreak on Obstetric Health Care in Sierra Leone. *PLoS One* 2016;11:e0150080.

7. Cancedda C, Davis SM, Dierberg KL, *et al*. Strengthening Health Systems While Responding to a Health Crisis: Lessons Learned by a Nongovernmental Organization During the Ebola Virus Disease Epidemic in Sierra Leone. *J Infect Dis* 2016; 214 (Suppl. 3):S153-S163.

8. Jones S, Sam B, Bull F, Pieh SB, *et al*. 'Even when you are afraid, you stay': Provision of maternity care during the Ebola virus epidemic: A qualitative study. *Midwifery* 2017;52:19-26.

9. Dynes MM, Miller L, Sam T, *et al*. Perceptions of the risk for Ebola and health facility use among health workers and pregnant and lactating women Kenema District, Sierra Leone, 2014. *Morb Mortal Wkly Rep* 2015;63:12267.

10. Delamou A, Hammonds RM, Caluwaerts S, *et al*. Ebola in Africa: beyond epidemics, reproductive health in crisis. *Lancet* 2014; 13; 384:2105.

11. Hayden EC. Ebola obstructs malaria control. *Nature* 2014;514:15-6.

12. Sierra Leone, Ministry of Health and Sanitation, United Nations Children’s Fund. *Health facility survey. Assessing the impact of the EVD outbreak on health systems in Sierra Leone*. MoHS, UNICEF, 2014.

13. Sierra Leone, Ministry of Health and Sanitation, UN Children's Fund. *Sierra Leone health facility assessment 2015: impact of the EVD [Ebola Virus Disease] outbreak on Sierra Leone’s primary health care system*. MoHS, UNICEF, 2015.

14. Jones S, Gopalakrishnan S, Ameh CA, *et al.* 'Women and babies are dying but not of Ebola': the effect of the Ebola virus epidemic on the availability, uptake and outcomes of maternal and newborn health services in Sierra Leone. *BMJ Glob Health* 2016;1(3):e000065.
15. Centers for Disease Control and Prevention (CDC). 2014-2016 Ebola Outbreak Distribution in West Africa. 2017. Available from: <https://www.cdc.gov/vhf/ebola/history/2014-2016-outbreak/distribution-map.html> [Accessed Dec 2018].
16. Sierra Leone, Ministry of Health and Sanitation. *Nationwide needs assessment for emergency obstetric and new-born care services in Sierra Leone*. MoHS, 2008.
17. Theuring S, Koroma AP, Harms G. "In the hospital, there will be nobody to pamper me": a qualitative assessment on barriers to facility-based delivery in post-Ebola Sierra Leone. *Reprod Health* 2018;15:155: 10.1186/s12978-018-0601-9.
18. Quaglio GL, Pizzol D, Bome D, *et al.* Maintaining maternal and child health services during the Ebola outbreak: experience from Pujehun, Sierra Leone. *PLoS Currents* 2016;8: ecurrents.outbreaks.d67aea257f572201f835772d7f188ba5.
19. Ly J, Sathananthan V, Griffiths T, *et al.* Facility-based delivery during the Ebola virus disease epidemic in rural Liberia: analysis from a cross-sectional, population-based household survey. *PLoS Med* 2016;13: e1002096.
20. Barden-O'Fallon J, Barry MA, Brodish P, *et al.* Rapid assessment of Ebola-related implications for reproductive, maternal, newborn, and child health service delivery and utilization in Guinea. *PLoS Curr* 2015;7:ecurrents.outbreaks.0b0ba06009d d091bc39ddb3c6d7b0826.
21. Lori JR, Rominski SD, Perosky JE, *et al.* A case series study on the effect of Ebola on facility-based deliveries in rural Liberia. *BMC Pregnancy Childbirth* 2015;15:254.
22. Iyengar P, Kerber K, Howe CJ, *et al.* Services for mothers and newborns during the Ebola outbreak in Liberia: the need for improvement in emergencies. *PLoS Curr* 2015; 7:ecurrents.outbreaks.4ba318308719ac86fbef91f8e56cb66f.
23. World Health Organization. *Situation report: Ebola virus disease, 10 June 2016*. WHO, 2018. Available from: [http://apps.who.int/iris/bitstream/10665/208883/1/ebolasitrep\\_10Jun2016\\_eng.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/208883/1/ebolasitrep_10Jun2016_eng.pdf?ua=1) [Accessed Dec 2018].
24. R Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. Vienna, Austria, 2018.
25. Camara BS, Delamou A, Diro E, *et al.* Effect of the 2014/2015 Ebola outbreak on reproductive health services in a rural district of Guinea: an ecological study. *Trans R Soc Trop Med Hyg* 2017;11:22-9.

26. Delamou A, Hammonds RM, Caluwaerts S, *et al.* Effect of Ebola virus disease on maternal and child health services in Guinea: a retrospective observational cohort study. *Lancet Glob Health* 2017;5:e448-e457.
27. Ajelli M, Parlamento S, Bome D, *et al.* The 2014 Ebola virus disease outbreak in Pujehun, Sierra Leone: epidemiology and impact of interventions. *BMC Med* 2015;13:281.
28. The Guardian. Sierra Leone declares first Ebola free district. 10 January 2015. Available from: <http://www.theguardian.com/world/2015/jan/10/sierraleonefirstebolafreedistrictwho> [Accessed Dec 2018].
29. Statistics Sierra Leone, United Nations Children's Fund. *Sierra Leone multiple indicator cluster survey 2017. Survey findings report.* Stats SL, UNICEF, 2018.
30. Sierra Leone, Ministry of Health and Sanitation. *Sierra Leone demographic and health survey.* MoHS, 2013.
31. Wagenaar BH, Augusto O, Beste J, *et al.* The 2014-2015 Ebola virus disease outbreak and primary healthcare delivery in Liberia: Time-series analyses for 2010-2016. *PLoS Med* 2018;15:e1002508.
32. Nobles J, Frankenberg E, Thomas D. The Effects of mortality on fertility: population dynamics after a natural disaster. *Demography* 2015;52:15–38.
33. Hapsari ED, Widyawati, Nisman WA, *et al.* Change in contraceptive methods following the Yogyakarta earthquake and its association with the prevalence of unplanned pregnancy. *Contraception* 2009;79:316-22.
34. McBain RK, Wickett E, Mugunga JC, *et al.* The post-Ebola baby boom: time to strengthen health systems *Lancet* 2016;388: 2331-3.
35. Bolkan HA, Bash-Taqi DA, Samai M, *et al.* Ebola and indirect effects on health service function in Sierra Leone. *PLoS Curr* 2014;6:ecurrents.outbreaks.0307d588d f619f9c9447f8ead5b72b2d.
36. Olu O, Kargbo B, Kamara S, *et al.* Epidemiology of Ebola virus disease transmission among health care workers in Sierra Leone, May to December 2014: a retrospective descriptive study. *BMC Infect Dis* 2015;15:416.
37. Thaddeus S, Maine D. Too far to walk: maternal mortality in context. *Sm Sci Med* 1994;38:1091-1110.
38. Treacy L. Distance, accessibility and costs. Decision-making during childbirth in rural Sierra Leone: A qualitative study. *PLoS One* 2018;13: e0188280.
39. Tayler-Smith K, Zachariah R, Manzi M, *et al.* An ambulance referral network improves access to emergency obstetric and neonatal care in a district of rural Burundi with high maternal mortality. *Trop Med Int Health* 2013;18:993-1001.

- 1  
2  
3 40. Groppi L, Somigliana E, Pisani V, *et al.* A hospital-centred approach to improve emergency  
4 obstetric care in South Sudan. *Int J Gynaecol Obstet* 2015;128:58–61.  
5  
6  
7 41. Tsegaye A, Somigliana E, Alemayehu T, *et al.* Ambulance referral for emergency obstetric care  
8 in remote settings. *Int J Gynaecol Obstet* 2016;133:316–9.  
9  
10  
11 42. Oyerinde K, Harding Y, Amara P, *et al.* Barriers to uptake of emergency obstetric and newborn  
12 care services in Sierra Leone: a qualitative study. *Comm Med & Health Educ* 2012;2:  
13 10.4172/2161-0711.1000149.  
14  
15 43. Vallières F, Cassidy EL, McAuliffe E, *et al.* Can Sierra Leone maintain the equitable delivery of  
16 their Free Health Care Initiative? The case for more contextualised interventions: results of a  
17 cross-sectional survey. *BMC Health Serv Res* 2016;16:258.  
18  
19  
20 44. Kruk ME, Myers M, Varpilah ST, *et al.* What is a resilient health system? Lessons from Ebola.  
21 *Lancet* 2015;385:1910–12.  
22  
23  
24 45. Witter S, Hunter B. *Resilience of health systems during and after crises – what does it mean and*  
25 *how can it be enhanced?* ReBUILD Research Programme Consortium. 2017. Available  
26 from:[https://rebuildconsortium.com/media/1535/rebuild\\_briefing\\_1\\_june\\_17\\_resilience.pdf](https://rebuildconsortium.com/media/1535/rebuild_briefing_1_june_17_resilience.pdf)  
27 [Accessed Dec 2018].  
28  
29  
30 46. Mussah VG, Mapleh L, Ade S, *et al.* Performance-based financing contributes to the resilience  
31 of health services affected by the Liberian Ebola outbreak. *Public Health Action* 2017;7(Suppl.  
32 1):S100–S105.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1 Pediatric and maternal admissions at hospital level




Figure 2 C-sections, deliveries, MDOCs, pediatric and maternal deaths at hospital level





Figure 3 ANC 1, ANC 4, deliveries, and family planning at community level





# BMJ Open

## Impact of Ebola outbreak on reproductive health services in a rural district of Sierra Leone. A prospective observational study.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-029093.R1
Article Type:	Research
Date Submitted by the Author:	21-Mar-2019
Complete List of Authors:	<p>Quaglio , GianLuca ; European Parliament  Tognon, Francesca; Department for Woman and Child Health, University of Padua, Italy  Finos, Livio; Department of Developmental Psychology and Socialisation, University of Padua, Italy  Bome, David; Ministry of Health and Sanitation, Freetown, Sierra Leone  Sesay, Santigie ; Ministry of Health and Sanitation, Freetown, Sierra Leone  Atiba, Kebbie ; Department for Woman and Child Health, Pujehun Hospital, Sierra Leone  Di Gennaro, Francesco ; Department of Infectious Diseases, University of Bari, Italy  Bienvenu Salim, Camara ; National Centre for Training and Research in Rural Health of Maferinyah, Forécariah, Guinea  Marotta, Claudia; Department of Science for Health Promotion and Mother to Child Care, University of Palermo, Italy  Pisani, Vincenzo; Department for Woman and Child Health, Pujehun Hospital, Sierra Leone, Pujehun, SL  Bangura , Zainab; Department for Woman and Child Health, Pujehun Hospital, Sierra Leone  Pizzol, Damiano; Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy  Saracino, Annalisa; Department of Infectious Diseases, University of Bari, Italy  MAZZUCCO, WALTER; Department of Science for Health Promotion and Mother to Child Care, University of Palermo, Italy  Jones, Susan; Department of Nursing and Midwifery School of Human and Health Sciences, University of Huddersfield, UK  Putoto, G; Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy</p>
<b>Primary Subject Heading</b>:	Health policy
Secondary Subject Heading:	Public health, Mental health, Infectious diseases, Addiction, Communication
Keywords:	PAEDIATRICS, Public health < INFECTIOUS DISEASES, Community gynaecology < GYNAECOLOGY



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



SCHOLARONE™  
Manuscripts

# Impact of Ebola outbreak on reproductive health services in a rural district of Sierra Leone. A prospective observational study.

Gianluca Quaglio,<sup>1,2,3</sup> Francesca Tognon,<sup>4</sup> Livio Finos,<sup>5</sup> David Bome,<sup>6</sup> Sesay Santigie,<sup>6</sup> Atiba Kebbie,<sup>7</sup> Francesco Di Gennaro,<sup>8</sup> Bienvenu Salim Camara,<sup>9</sup> Claudia Marotta,<sup>10</sup> Vincenzo Pisani,<sup>7</sup> Bangura Zainab,<sup>7</sup> Damiano Pizzol,<sup>3</sup> Annalisa Saracino,<sup>8</sup> Walter Mazzucco,<sup>10</sup> Susan Jones,<sup>11</sup> Giovanni Putoto<sup>3</sup>

1. European Parliamentary Research Services (EPRS), European Parliament, Brussels, Belgium;
2. Department of International Health/CAPHRI, Maastricht University, The Netherlands;
3. Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy;
4. Department for Woman and Child Health, University of Padua, Italy;
5. Department of Developmental Psychology and Socialisation, University of Padua, Italy;
6. Ministry of Health and Sanitation, Freetown, Sierra Leone;
7. Department for Woman and Child Health, Pujehun Hospital, Sierra Leone;
8. Department of Infectious Diseases, University of Bari, Italy;
9. National Centre for Training and Research in Rural Health of Maferinyah, Forécariah, Guinea;
10. Department of Science for Health Promotion and Mother to Child Care, University of Palermo, Italy;
11. Department of Nursing and Midwifery School of Human and Health Sciences, University of Huddersfield, UK.

## Corresponding author

Gianluca Quaglio, MD, PhD  
Directorate-General for Parliamentary Research Services (EPRS)  
European Parliament  
Rue Wiertz, 60, B-1047  
Brussels, Belgium  
gianluca.quaglio@europarl.europa.eu

**Words count** (excluding title page, abstract, references, figures and tables): 5182



1

2

3 47

4 48

5

6 49

7

8 50

9

10 51

11

12 52

13

14 53

15 54

16

17 55

18

19 56

20

21 57

22

23 58

24

25 59

26 60

27

28 61

29

30 62

31

32 63

33

34 64

35

36 65

37 66

38

39 67

40

41 68

42

43 69

44

45 70

46

47 71

48 72

49

50 73

51

52 74

53

54 75

55

56 76

57 77

58

59 78

60 79

**ABSTRACT**

**Objectives** To assess the trends concerning utilisation of maternal and child health (MCH) services before, during, and after the Ebola outbreak, quantifying the contribution of a reorganised referral system (RS).

**Design** A prospective observational study of MCH services.

**Setting** Pujehun district in Sierra Leone, 77 community health facilities and 1 hospital from 2012 to 2017.

**Main outcome measures** MCH utilization was evaluated by assessing: i) institutional deliveries, Cesarean-sections, paediatric and maternity admissions and deaths, and major direct obstetric complications (MDOCs), at hospital level; ii) antenatal care (ANC) 1 and 4, institutional delivery, and family planning, at community level. Contribution of a strengthened RS was also measured.

**Results** At hospital level, there is a significant difference between trends Ebola vs pre-Ebola for maternal admissions (7, 95% CI 4 to 11, p <0.001), MDOCs (4, 95% CI 1 to 7, p = 0.006), and institutional deliveries (4, 95% CI 2 to 6, p = 0.001). There is also a negative trend in the transition from Ebola to post Ebola for maternal admissions (-7, 95% CI -10 to -4, p <0.001), MDOCs (-4, 95% CI -7 to -1, p 0.009) and institutional deliveries (-3, 95% CI -5 to -1, p 0.001). The differences between trends pre-Ebola vs post-Ebola are only significant for pediatric admissions (3, 95% CI 0 to 5, p 0.035). At community level, the difference between trends Ebola vs pre-Ebola and Ebola vs post-Ebola are not significant for any indicators. The differences between trends pre-Ebola vs post-Ebola show a negative difference for institutional deliveries (-7, 95% CI -10 to -4, p <0.001) ANC 1 (-6, 95% CI -10 to -3, p <0.001), ANC 4 (-8, 95% CI -11 to -5, p <0.001) and family planning (-85, 95% CI -119 to -51, p <0.001).

**Conclusions** A stronger health system compared to other districts in Sierra Leone and a strengthened RS enabled health facilities in Pujehun to maintain service provision and uptake during and after the Ebola epidemic.

**Keywords:** Ebola, Sierra Leone, Maternal and Child Health indicators, Referral system, Reproductive health service.

## Strengths and limitations of this study

- ▶ The study uses data from a remote rural district in Sierra Leone, with a 6-year observational period. Data have been collected in a prospective way, reducing the potential bias in the accuracy of the data reported by other studies carried out in countries affected by Ebola.
- ▶ Data from pre, intra, and post-Ebola periods allowed comparisons between trends, something rarely carried out in countries heavily affected by Ebola.
- ▶ The data refers to a single area of Sierra Leone: the sample cannot be considered representative of the country as a whole.
- ▶ In addition to measures put in place to reduce the impact of the disease on mothers and children, Pujehun had far fewer Ebola cases than other districts, which may also have led to the utilization of health services.

1

2

3 92 INTRODUCTION

4

5 93 The 2014-2015 Ebola Virus Disease (EVD) outbreak was the most severe in history, mainly affecting

6

7 94 three West African countries; Guinea, Sierra Leone and Liberia. Overall 28,616 people were infected

8

9 95 of which 11,310 died and the outbreak was declared a global public health emergency by the WHO.<sup>1</sup>

10

11 96 Of the three countries affected, Sierra Leone had the most confirmed cases (8,704), which accounted

12

13 97 for 50% of all confirmed cases in West Africa, and 3,589 deaths.<sup>2-4</sup> All 14 districts in Sierra Leone were

14

15 98 affected, but at different times and to varying degrees.<sup>5</sup> During the Ebola crisis the population's trust

16

17 99 in the national health system declined in Sierra Leone, leading to an overall reduction in the use of

18

19 100 health services, including reproductive, maternal, and child services.<sup>6-8</sup> Underlying factors for the

20

21 101 decrease in the use of health services included fear of infection, for both healthcare workers and

22

23 102 patients, the underlying fragility of the health systems, the reduced numbers of available health

24

25 103 personnel, and the death of healthcare workers due to EVD.<sup>9 10</sup> It has been estimated that 30% of

26

27 104 health workers who died of EVD in West Africa were maternal and child healthcare (MCH) providers.<sup>11</sup>

28

29 105 However, there were considerable variations in the reduction of health service uptake when looked

30

31 106 at by district level in Sierra Leone.<sup>6 12-14</sup> While districts such as Kambia, Port Loko and Bonthe showed

32

33 107 large reductions in facility-based delivery (between 38-41%), the district of Pujehun showed only a

34

35 108 5% decrease in the same service. Similar geographic variations were seen in the reduction in antenatal

36

37 109 care (ANC) visits.<sup>12 13</sup>

38

39 110 The number of confirmed EVD cases - and deaths - varied considerably by district. There were

40

41 111 no more than 100 confirmed cases in both Bonthe and Pujehun, and up to 4,000 confirmed cases in

42

43 112 both Port Loko and Bombali.<sup>15</sup> However, public fear of Ebola, regardless of the actual number of cases

44

45 113 per district, may still have prevented many people from accessing services. The challenge of providing

46

47 114 adequate levels of care during a humanitarian emergency such as the EVD crisis was further

48

49 115 exacerbated by the weak health system in Sierra Leone, particularly in rural areas where the poor

50

51 116 condition of the roads and high transport costs cause delays in accessing services, and contribute to

52

53 117 increased maternal and neonatal mortality.<sup>16</sup>

54

55 118 Doctors with Africa (DwA) CUAMM is an Italian NGO working in Sierra Leone since 2012. It is

56

57 119 present in the Pujehun district focusing on MCH care both at hospital and community level<sup>17 18</sup> In this

58

59 120 paper, community level refers to Peripheral Health Units (PHUs), i.e. all health facilities outside the

60

121 hospital. As described in our previous reports,<sup>17 18</sup> a number of measures were put in place to control

122 the Ebola epidemic in the Pujehun district which reduced the impact of the disease on mothers and

123 children compared to other districts. During this EVD epidemic, the predominantly vertical focus on

1  
2  
3 124 outbreak control was associated with failures in providing effective care for routine health needs.<sup>19-</sup>  
4  
5 125 <sup>21</sup> In contrast, the approach implemented in the Pujehun district was not based on vertical actions  
6  
7 126 and 'humanitarian response to health emergencies with a short half-life'.<sup>21</sup> Rather, it worked on  
8  
9 127 strengthening all the components of the health system - governance, human resources, community  
10  
11 128 involvement - before, during and, after the epidemic. A rapid response to the crisis by the local health  
12  
13 129 authorities was implemented adopting public health measures before any other district in Sierra  
14  
15 130 Leone.<sup>22</sup> The activities were mainly concentrated on keeping the health service open and properly  
16  
17 131 functioning in order to reduce the collateral effects of the epidemic on routine health services. No  
18  
19 132 health units in the Pujehun district were closed during the epidemic. Measures to empower  
20  
21 133 community leaders and use culturally appropriate methods of communication helped to dispel  
22  
23 134 community mistrust in the health services. At community level, a number of strategies were  
24  
25 135 implemented such as the regular rotation of health facility staff, which strengthened teamwork and  
26  
27 136 effective leadership. In Sierra Leone, healthcare workers based at community health centres may  
28  
29 137 often work alone in isolated centres with limited support from clinical colleagues or management. By  
30  
31 138 rotating staff through the various facilities, they gain on the job training, peer support, and develop  
32  
33 139 new working relationships. At the start of the Ebola epidemic, many expatriate healthcare workers  
34  
35 140 in NGOs left Sierra Leone, negatively affecting care delivery and staff morale. The continued presence  
36  
37 141 of international teams in the daily activities in Pujehun hospital and the acceptance of the  
38  
39 142 professional risks by both national and international staff may have contributed to maintaining an  
40  
41 143 attitude of 'normality' in an extremely stressful environment. This might also help to explain the  
42  
43 144 population's positive receptiveness towards the health services.<sup>17 18</sup>

44  
45 145 Different types of referral systems (RSs) such as motorbikes were present in the country in the  
46  
47 146 pre Ebola period to transport patients from the villages to the nearest health facility. Ambulances  
48  
49 147 were also present in several districts with 73% of health facilities nationwide having a functioning RS,  
50  
51 148 59% of them consisting of an ambulance on call.<sup>12 23</sup> In the Pujehun district, the RS was barely  
52  
53 149 functioning, only able to support the activity of a limited number of PHUs. The service was also  
54  
55 150 entirely funded by the patients themselves, resulting in underutilization of the service. Utilization was  
56  
57 151 further reduced during the outbreak, when the ambulances were identified by the population with  
58  
59 152 the transport of Ebola infected patients, and their use occasioned fear and distrust. In January 2015,  
60  
61 153 in collaboration with the Ministry of Health and Sanitation (MoHS) of Sierra Leone and UNICEF, DWA  
62  
63 154 began the re-organisation and reinforcement of the RS, transferring pregnant women and pediatric  
64  
65 155 cases from PHUs to the Pujehun hospital.

Our previous studies<sup>18</sup> provided information only on three MCH indicators, namely pediatric admissions, maternity admissions, and institutional deliveries; in addition it did not assess the trends in the post-EVD period. Existing studies examining the influence of EVD on MCH services targeted the outbreak and the immediate post-outbreak periods.<sup>24-27</sup> Understanding the trends in the use of MCH services before, during, and after the EVD outbreak will help to guide post-EVD interventions, increasing access to MCH services in rural Sierra Leone. This information will also be useful in preparing a more organised and structured RS. With this background, the aims of this study are: i) to assess trends in institutional deliveries, C-sections, paediatric and maternity admissions, paediatric and maternity deaths, and major direct obstetric complications (MDOCs), before, during, and after the EVD in the Pujehun hospital, thus complementing the results of the previous report which were limited to 3 MCH indicators; ii) to assess trends in ANC 1 and 4, institutional delivery, and family planning, at community level. This study was carried out in conjunction with the strengthening of an RS initiated a few weeks after the Pujehun district was declared Ebola-free.

**METHODS**

**Setting**

Sierra Leone has four provinces that are divided into 14 districts. Pujehun is one of four districts in the southern province (Figure 1). It has a population of approximately 375,000 inhabitants. The primary care network included 77 MoHS PHUs, 5 of which provide basic emergency obstetrics care (BEmOC). The secondary care system consists of the MoHS provided district hospital, which comprises the MCH complex, providing comprehensive emergency obstetric and newborn care (CEmONC) services. Connections between the community and health facilities are difficult because of the very poor condition of the roads. Furthermore, the district is divided by a major river (Moa River) and has a riverine area reachable only with boats, which further hinders access. The first case of Ebola in Pujehun district was reported on the 7th July 2014. The district was declared Ebola free on the 10th January 2015.<sup>28</sup> A total of 49 patients were registered with a case fatality rate of 85.7% (42/49).

**Referral system**

In the Pujehun district, two ambulances managed by the District Health Management Team (DHMT) were functioning in the pre Ebola period, but only 63% of the PHUs were able to use the service.<sup>12 23</sup> Emergency calls were not coordinated by the hospital and the transport costs were covered by the patients, dissuading many from using the service. During the outbreak, people came to associate the

ambulances with transporting Ebola infected patients, which further discouraged their use. A 24-h free-of-charge ambulance RS, transferring pregnant women with obstetric complications from the health centers to Pujehun hospital was implemented in January 2015. In the hospital a call center was established and the call center number was distributed to all the 77 PHUs. Private calls were considered only in the case of an emergency or if the staff of the PHU were not available. After confirming an emergency condition together with the PHU staff, the hospital midwife had the responsibility to authorize the referral. A nurse on duty from the maternity hospital accompanied the driver in each referral. Health personnel at hospital and PHUs levels were trained on Life Saving Skills – Emergency Obstetric and Newborn Care, including referral criteria and definition of MDOCs.<sup>29</sup>

Referrals were carried out by 3 ambulances, two positioned in the Pujehun MCH complex, and a third one in Jendema, bordering Liberia, on the opposite side of the Moa River. Around the Jendema area, 15 PHUs were located serving a population of approximately 80,000 inhabitants. Referrals in this area were made using the ambulances and by transferring patients at the river crossing point via a barge or a motor boat, depending on the flow rate of the river. Pediatric referrals were performed using private motor bikes available in the villages and hired from PHUs staff without the involvement of the call center. A referral form describing the clinical case and the justification for the referral was distributed to all the PHUs. The bike rider, after bringing the patient to the pediatric ward, delivered the referral form and received the reimbursement. For all patients carried to the hospital information was collected, including demographics, location, and the reason for contacting the RS. Community awareness activities were organized about the RS through meetings and radio discussions held by the DMHT, hospital health personnel, and local authorities.

### **Study design, population, and period**

A prospective observational study using routinely collected health services data, from January 2012 to December 2017, was carried out. Three time periods were considered: pre- Ebola period (1<sup>st</sup> January 2012 – 30<sup>th</sup> May 2014); Ebola period (1<sup>st</sup> June 2014 – 28<sup>th</sup> February 2015); post- Ebola period (1<sup>st</sup> March 2015 – 31<sup>th</sup> December 2017). We considered the Ebola period from one month before the first confirmed case in the district (i.e. June 2014), to one month after the country being declared Ebola free (i.e. February 2015). This was done because in Sierra Leone the outbreak had started in other districts of the country before the first case registered in Pujehun and continued to affect other districts until November 2015. It is realistic to assume that public fear of potential EVD cases and lack of confidence in the health services persisted in the Pujehun population during that time.<sup>14</sup> In



1  
2  
3 220 addition, expanding the Ebola period enabled a full assessment of the impact of the disease with an  
4  
5 221 adequate comparison with the two long periods before and after the Ebola epidemic.  
6  
7 222

8  
9 223 **Data collection**

10 224 Data on MCH indicators was prospectively collected from hospital registers (maternity ward, delivery  
11  
12 225 unit, pediatric ward, operating theatre). The following variables were collected on a monthly basis:  
13  
14 226 1) paediatrics admissions; 2) pediatric deaths; 3) maternity admissions; 4) maternal deaths; 5)  
15  
16 227 deliveries; 6) C-sections; 7) MDOC cases. MDOC cases were collected using a dedicated database  
17  
18 228 within the hospital and confirmed by a gynaecologist. All hospital maternal deaths were reviewed by  
19  
20 229 DHMT and classified according to Maternal Death Surveillance and Response policy by MoHS.  
21  
22 230 Paediatric deaths did not include stillbirths and early neonatal deaths, but only deaths of children  
23  
24 231 admitted to the paediatric ward.

25 232 At community level, the following variables were collected from the local district Health  
26  
27 233 Management Information System (HMIS): 1) family planning consultations per month; 2) deliveries  
28  
29 234 per month; 3) ANC 1 per month; 4) ANC 4 per month. Different variables were collected from the  
30  
31 235 two types of sites, based on the different services provided at community level (BEmOC) and at  
32  
33 236 hospital level (CEmONC). Quarterly review meetings were organized with the staff in charge of the  
34  
35 237 health facilities to address data discrepancies in the reports. Technical assistance was provided to  
36  
37 238 the DHMT to improve timeliness, completeness, and accuracy of data regarding CEmOC and BEmONC  
38  
39 239 services.

40 240 For the RS, data was collected from records of all of the study sites, including delivery registers,  
41  
42 241 delivery logbooks, prenatal registers, referral registers, and death registers. Additional data was  
43  
44 242 collected from the ambulance database and logbook. Records in the database were then validated  
45  
46 243 by cross-checking the records with registers at the study sites.  
47  
48 244

49 245 **Statistical analysis**

50  
51 246 For each indicator, a segmented seasonal autoregressive model of order 1 was estimated. The  
52  
53 247 segments defined the three periods: before the EVD epidemic (January 2012 to May 2014), during  
54  
55 248 the epidemic (June 2014 to February 2015), and after the epidemic (March 2015 to December 2017).  
56  
57 249 The model for each indicator  $Y_t$  collected at hospital or community level was as follows:  $Y_t = \beta_0 + \beta_1$   
58  
59 250  $T_t + \beta_2 X_t + \beta_3 X_t T_t + \beta_4 Z_t + \beta_5 Z_t T_t + \beta_6 Month + \varepsilon_t$ .  $\beta_0$  estimates the number of individuals using  
60  
61 251 the service at the beginning of the pre-Ebola period;  $\beta_1$  estimates the average monthly change in the

number using the service over the pre-outbreak period;  $T_t$  is the time since the start of the study;  $\beta_2$  represents the change in the level of service use that occurred in the period immediately after the EVD period (designated by indicator variable  $X_t$ );  $\beta_3$  represents the difference between the trend in service use during the EVD outbreak compared to the pre-disease period;  $\beta_4$  represents the change in service use that occurred in the period immediately after the end of the outbreak (post-outbreak period designated by indicator variable  $Z_t$ );  $\beta_5$  is the difference between the trend in service use during the period after the Ebola virus disease outbreak compared with the period during the outbreak period;  $\beta_m$  represents a series of indicator variables for each calendar month, and  $t$  is the random error term. Overall trends across the periods and the comparisons among trends were calculated as follows: linear trend during the outbreak =  $\beta_1 + \beta_3$ ; linear trend after the outbreak =  $\beta_1 + \beta_3 + \beta_5$ ; and linear trend after the outbreak vs linear trend before the outbreak =  $\beta_3 + \beta_5$ . Average levels across the periods and their comparisons were calculated as follows: average during the outbreak =  $\beta_0 + \beta_2$ ; average after the outbreak =  $\beta_0 + \beta_2 + \beta_4$ ; and difference between after the outbreak and before the outbreak =  $\beta_2 + \beta_4$ . Differences were considered statistically significant at  $p < 0.05$ . The analysis was performed using R.<sup>30</sup> The full data analysis is available in Annex 1.

### Patient involvement

No patients were involved in defining the research question or the outcome measures, nor were they involved in the design and implementation of the study. There are no plans to involve patients in the dissemination of the results.

## RESULTS

### Hospital level: Pre-Ebola period

At hospital level, for all indicators, the trend is stable during the pre Ebola period, without significant changes (Figure 2 and 3).

### Hospital level: Ebola vs pre-Ebola period

At hospital level, the differences between Ebola period vs pre-Ebola averages show a statistically significant increase for institutional deliveries (11, 95% CI 2 to 21,  $p = 0.02$ ) and for the reduction of maternal deaths (-1, 95% CI - 2 to 0,  $p = 0.042$ ) (Table 1). There is also a statistically significant difference between the trend of Ebola period vs pre-Ebola period, for maternal admissions (7, 95%

1  
2  
3 283 CI 4 to 11, p <0.001), MDOCs (4, 95% CI 1 to 7, p = 0.006), and institutional deliveries (4, 95% CI 2 to  
4  
5 284 6, p = 0.001) (Figure 2 and 3).  
6  
7 285  
8  
9 286 **Hospital level: Ebola vs post-Ebola period**  
10  
11 287 At hospital level, the differences between averages of the post Ebola vs Ebola are statistically  
12  
13 288 significant for all indicators: institutional deliveries, C-sections, paediatric and maternity admissions,  
14  
15 289 paediatric and maternity deaths, and MDOCs (Table 1). There is also a negative trend in the transition  
16  
17 290 from Ebola to post Ebola for maternal admissions (-7, 95% CI -10 to -4, p <0.001), MDOCs (-4, 95% CI  
18  
19 291 -7 to -1, p 0.009) and institutional deliveries (-3, 95% CI -5 to -1, p 0.001) (Figure 2 and 3).  
20  
21 292  
22 293 **Hospital level: Pre-Ebola vs post-Ebola period**  
23  
24 294 The differences between averages of the pre-Ebola vs post-Ebola periods are also statistically  
25  
26 295 significant for all indicators, except for maternal deaths (Table 1). The differences between trends  
27  
28 296 between pre-Ebola vs post-Ebola period are only significant for pediatric admissions (3, 95% CI 0 to  
29  
30 297 5, p 0.035) (Figure 2 and 3).  
31  
32 298

Table 1 MCH indicators at hospital and community level									
Indicator	Difference between average of Ebola period vs pre-Ebola period			Difference between average of Ebola period vs post-Ebola period			Difference between average of pre-Ebola period vs post-Ebola period		
	β	95% CI	p value	β	95% CI	p value	β	95% CI	p value
<b>HOSPITAL LEVEL</b>									
Maternal admissions	7	-7 to 22	0.333	43	28 to 58	<0.001	50	37 to 64	<0.001
Maternal deaths	-1	-2 to 0	0.042	2	1 to 3	0.001	1	0 to 2	0.135
Institutional deliveries	11	2 to 21	0.02	28	18 to 38	<0.001	39	31 to 48	<0.001
C-sections	5	-1 to 11	0.13	15	8 to 21	<0.001	19	13 to 25	<0.001
MDOC	2	-11 to 14	0.782	41	30 to 54	<0.001	43	31 to 54	<0.001
Pediatric admissions	1	-39 to 40	0.968	133	92 to 174	<0.001	134	98 - 170	<0.001
Pediatric deaths	-1	-6 to 5	0.826	9	3 to 15	0.004	8	3 to 14	0.003
<b>COMMUNITY LEVEL</b>									
Institutional deliveries	148	99 to 196	<0.001	-10	-59 to 39	0.695	138	93 to 183	<0.001
ANC 1	74	3 to 145	0.042	-48	-122 to 26	0.2	26	-40 to 91	0.448
ANC 4	80	21 to 139	0.008	23	-38 to 84	0.461	103	48 to 157	<0.001
Family planning	490	-92 to 1073	0.099	-262	-855 to 330	0.386	228	-293 to 750	0.391

51 299  
52 300  
53 301  
54 302  
55 303  
56 304  
57  
58 303  
59  
60 304

### Community level: Pre-Ebola period

At community level, all indicators in the months before Ebola showed a positive trend. There was a monthly average increase of 8 institutional deliveries (95% CI 6 to 10,  $p < 0.001$ ); a monthly average increase of 7 ANC 1 (95% CI 4 to 10,  $p < 0.001$ ) and 6 ANC 4 (95% CI 4 to 8,  $p < 0.001$ ), and a monthly average increase of 69 women accessing family planning services (95% CI 42 to 95,  $p < 0.001$ ) (Figure 4).

### Community level: Ebola vs pre-Ebola period

At community level, with the exception of family planning, the differences between averages of Ebola period vs pre-Ebola are statistically significant for all indicators: institutional deliveries (148, 95% CI 99 to 196,  $p < 0.001$ ), ANC 1 (74, 95% CI 3 to 145,  $p = 0.042$ ), and ANC 4 (80, 95% CI 21 to 139,  $p = 0.008$ ) (Table 1). The difference between trends (Figure 3) of the Ebola vs pre-Ebola period are not significant for any of the indicators considered (Figure 4).

### Community level: Ebola vs post-Ebola period

At community level, the differences between averages (Table 1) and the difference between trends (Figure 4) of the Ebola vs post-Ebola period are not significant for any of the indicators considered.

### Community level: Pre-Ebola vs post-Ebola period

The differences between averages of the pre-Ebola vs post-Ebola are statistically significant, with an increase in institutional deliveries (138, 95% CI 93 to 183,  $p < 0.001$ ) and ANC 4 (103, 95% CI 48 to 157,  $p < 0.001$ ) (Table 1). However, there is a negative difference between trends among the two periods, for all the variables considered: institutional deliveries (-7, 95% CI -10 to -4,  $p < 0.001$ ) ANC 1 (-6, 95% CI -10 to -3,  $p < 0.001$ ), ANC 4 (-8, 95% CI -11 to -5,  $p < 0.001$ ) and most significantly for family planning (-85, 95% CI -119 to -51,  $p < 0.001$ ) (Figure 4).

### Referral system: Obstetric and paediatric results

Between January 2015 and December 2017 there were 2,450 obstetric referrals. Of these, 1,574 (64%) were MDOC, which represent 70% of all the 2,233 MDOCs treated in the hospital over the same period. The baseline characteristics and reasons for MDOCs collected through the RS are reported on Table 2. At the same time, 4,671 paediatric patients were admitted in the hospital through the RS,

1  
2  
3 337 representing 72% of the 6,518 total admission during the same period. Reasons for paediatric  
4  
5 338 referrals are shown on Table 3.

6  
7 339

8 340

9 341

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41 342

42 343

43 344

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 345

59 346

60 347

348

Table 2 Baseline characteristics and reasons for MDOCs collected through RS, period 2015 - 2017		
Age (years)	N	%
Mean	25,3	SD 7
12-19	442	28%
20-29	613	39%
30-39	464	29%
40+	43	3%
Unknown	12	1%
Number of previous deliveries		
0	474	30%
1 or 2	377	24%
3 or 4	292	19%
5 or 6	207	13%
7+	212	13%
Unknown	12	1%
MDOC treated		
Prolonged/obstructive labour	848	54%
Antepartum haemorrhage	195	12%
Severe pre-eclampsia/eclampsia	165	11%
Abortium complicatium	117	7%
Post-partum haemorrhage	157	10%
Ectopic pregnancy	24	2%
Rupture uterus	30	2%
Sepsis	38	2%
Total	1574	100%

Table 3 Reasons for paediatric RS, period 2015-2017*		
Reason for referral	Number	%
Malaria	1540	30%
Anemia	910	18%
Pneumonia/ARI**	830	16%
Diarrhoea and vomiting	495	10%
Malnutrition	274	5%
Convulsion	186	4%
Hernia/Hydrocele	165	3%
Sepsis/Septicemia	127	2%
Dehydration	48	1%
Burn	30	1%
Others	522	10%
Total	5127	100%

\* For a number of patients, more than one suspected diagnosis for referral was reported; \*\* Acute Respiratory Infection.

## DISCUSSION

This study presents for the first time trends in utilization of MCH services before, during, and after Ebola, at hospital and community level from the country most heavily affected by the Ebola epidemic. It also presents data on the restructured and reorganised RS, which started immediately after the EVD outbreak. The study shows that there was a decrease in all MCH indicators and service uptake immediately after the onset of the outbreak, with a levelling or increase during the EVD period. In the post-Ebola period, all indicators (except for maternal deaths) showed an increase, in comparison with the pre-Ebola period. This was particularly marked at hospital level because the post Ebola reinforcement of the RS led to an increase in pediatric admissions, maternal admissions, and consequently a rise of institutional deliveries, C-sections, and MDOCs. In addition, while at the hospital level trends in the post-Ebola period are in line with the pre-Ebola, at community level there is a negative trend compared to the pre-Ebola period for all indicators taken into consideration. The study presents results in contrast to other studies that showed a decline in MCH services in the Ebola and post-Ebola periods.<sup>6 31 32</sup>

### Pre Ebola and Ebola periods

As mentioned above, the approach implemented in the Pujehun district<sup>17 28</sup> avoided vertical interventions only focused on the containment of the EVD epidemic. It worked on strengthening all the components of the health system - before, during, and long after the epidemic. This approach reduced the spread of infection and the impact of the disease on MCH services.<sup>17 18</sup> As shown by this paper, at community level family planning, ANC, and institutional deliveries, were affected only at the beginning of the Ebola outbreak with a small decrease in service utilization. In contrast, Jones et al., evaluated the number of antenatal and postnatal visits, institutional births, emergency obstetric care (EmOC), maternal deaths and stillbirths across 13 districts of Sierra Leone for 10 months during, and 12 months prior to the epidemic. They found that following the onset of the epidemic there was an 18% decrease in the number of women attending ANC visits and an 11% decrease in the number of women attending for birth at healthcare facilities.<sup>14</sup>

During the Ebola epidemic, the Pujehun hospital maintained C-sections and delivery volume at pre-Ebola levels. There was a stable number of patients attending the hospital during the Ebola outbreak, as shown by the number of maternal and pediatric admissions. The study of Brolin and colleagues focused on in-hospital deliveries and C-section volume in Sierra Leone. They showed that nationwide, albeit with substantial variation between districts, in-hospital deliveries and C-sections

1  
2  
3 382 decreased by over 20% during the Ebola outbreak, mainly because of the closure of not-for-profit  
4  
5 383 hospitals.<sup>6</sup> Brolin also noted that in general, at hospital level, in Sierra Leone those facilities that  
6  
7 384 remained open performed about the same number of deliveries and C-sections after the onset of the  
8  
9 385 EVD outbreak as they did before.<sup>6</sup> This seems to indicate that the decrease observed at national level  
10  
11 386 was related to the closing of key health facilities. The number of Ebola cases was not uniform  
12  
13 387 throughout districts in Sierra Leone and Pujehun was one of the least affected districts. The low  
14  
15 388 number of cases may also have helped to maintain public confidence in service provision and uptake  
16  
17 389 of services.<sup>7 8</sup>

18  
19  
20 391 **Post Ebola period**

21  
22 392 There is a shortage of data in Sierra Leone and the other West Africa countries affected regarding the  
23  
24 393 resumption of services after the epidemic. Pujehun district showed contrasting results at community  
25  
26 394 level. Results of the post Ebola vs pre-Ebola show an increase of activities for institutional delivery  
27  
28 395 and ANC 4. However, there is a negative trend among the two periods, for the variables taken into  
29  
30 396 consideration, namely institutional deliveries (-7, 95% CI -10 to -4, p <0.001) ANC 1 (-6, 95% CI -10 to  
31  
32 397 -3, p <0.001), ANC 4 (-8, 95% CI -11 to -5, p <0.001) and family planning (-85, 95% CI -119 to -51, p  
33  
34 398 <0.001). In 2017, the Pujehun district showed a coverage of 98% for ANC 1 (98% in 2013), 91% for  
35  
36 399 ANC 4 (76% at national level in 2013) and 90% for institutional deliveries (62% in 2013).<sup>33 34</sup> The initial  
37  
38 400 intervention carried out by DWA in the period 2012-2014 at the community level probably increased  
39  
40 401 these percentages, with an initial growth of the trend that had been slowing down in the years 2016-  
41  
42 402 2017. Possible explanations for this may include: bypassing, i.e. using alternative health care instead  
43  
44 403 of free or subsidized public clinics; increased opportunities to get transport to seek healthcare in  
45  
46 404 neighbouring districts; reduced demand for MCH services at community level; and reduced quality of  
47  
48 405 MCH services at PHUs.

49  
50 406 A study by Camara et al. in a rural district of Guinea showed a considerable recovery gap in the  
51  
52 407 post-Ebola period for ANC (37%) and institutional deliveries (34%).<sup>31</sup> Also Delamou et al. noted a  
53  
54 408 significant reduction in the average number of ANC visits and institutional deliveries during the Ebola  
55  
56 409 outbreak, in 6 districts of Guinea, and the overall post-outbreak trends did not suggest recovery.<sup>32</sup> By  
57  
58 410 contrast, Wagenaar et al., which analysed 10 primary care indicators in Liberia, before, during, and  
59  
60 411 after the Ebola outbreak, showed significant positive trends during the post-EVD period for ANC and  
61  
62 412 institutional deliveries.<sup>35</sup>



There are multifactorial and complex reasons for the decline of family planning in the Pujehun district. The activities that MoHS and DWA implemented from 2012 onwards were maintained during and after the EVD epidemic. However, a general decrease in the availability of healthcare personnel and international aid was observed and this could be a factor in the family planning decline. A possible stock-out of family planning methods has also been suggested as a reason for the decrease.<sup>25</sup> In addition, a reduction in demand for family planning in the post Ebola period could account for the decline of the service. Experiencing a disaster can trigger the desire to “rebuild” communities, reducing the need for family planning methods,<sup>36</sup> or communities may prefer traditional methods of contraception.<sup>37</sup> However, the reduction in family planning use in Pujehun district did not translate into an increase in institutional deliveries as occurred in neighbouring Liberia.<sup>38</sup> Although no further transmissions of Ebola took place in the Pujehun district after November 2015, the awareness of the ongoing transmission elsewhere in Sierra Leone, in Guinea and Liberia might have influenced health seeking behaviours.<sup>39 40</sup> However, this does not seem to have influenced other types of MCH services at community level. For comparison, the above mentioned study of Camara et al. showed that the utilization of family planning declined by 51% during the Ebola outbreak but recovered in the post-Ebola period.<sup>31</sup>

At hospital level, the situation is different. In the post-Ebola period, there was a significant increase in the volumes of activities: pediatric and maternal admissions, MDOC cases, deliveries, and C-sections. This increase can be directly linked to the reorganization and strengthening of the RS immediately after the Ebola epidemic. Based on the 3 delays theory,<sup>41</sup> in Pujehun it was decided to tackle the second delay, a lack of accessibility to health services. The distance to the hospital as well as lack of accessible and affordable vehicles were recognized as significant barriers when attempting to access CEmONC services at the hospital.<sup>42 43</sup> The success of the RS service can be linked to the integration of the key components needed for a successful service, namely: i) a transport system which took account of the specific geographical characteristics of the district;<sup>42</sup> ii) an effective communication system with a call center in contact with all PHUs of the district, the ambulance drivers, and the hospital; iii) training of all the PHU staff on the recognition of obstetric emergencies and on the RS.<sup>44 45</sup> Several meetings were planned with local community leaders and religious leaders to raise awareness of the importance of giving birth in health facilities. Prohibitive costs have been shown to be a major factor in preventing women accessing health facilities during childbirth in Sierra Leone.<sup>42 46 47</sup> Meetings were also organised to inform the population that the service was free of charge, and to give reassurance that the ambulances carried no risk of Ebola infection to people using



1  
2  
3 445 them. The increase in complicated cases treated at the hospital did not translate into an increase in  
4  
5 446 maternal and pediatric deaths, reflecting positively on the quality of care provided. The maternity  
6  
7 447 ward death rate remained around 1% throughout the 2012-2017 study period. The differences in  
8  
9 448 average death rates during the period 2015-2017 among referred and not referred pediatric patients  
10  
11 449 were 10.5% and 4.3% respectively. This showed that the pediatric RS works for the most critical cases  
12  
13 450 able to reach the hospital in time.

14 451  
15  
16 452 **CONCLUSIONS**

17  
18 453 There are a number of contextual factors and limitations that should be taken into account in the  
19  
20 454 analysis of the results of this study. The data refers to a single area of Sierra Leone and therefore our  
21  
22 455 sample cannot be considered representative of the country as a whole. We defined our distinct period  
23  
24 456 of EVD outbreak arbitrarily, from one month before the first case in the district to three months after  
25  
26 457 the last case in the district. This was done because the EVD crisis affected areas of the country outside  
27  
28 458 Pujehun prior to and after outbreak within Pujehun. The official end of the EVD epidemic for Sierra  
29  
30 459 Leone was declared on March 17, 2016, and for the countries of Guinea and Liberia was declared on  
31  
32 460 June 1, 2016. Finally, our study assumed that no other interventions in addition to those described  
33  
34 461 occurred concurrently with the Ebola epidemic.<sup>18</sup> Similarly, we assumed that no other substantial  
35  
36 462 interventions in addition to the re-organisation of the RS happened in the post-Ebola period which  
37  
38 463 would have affected the service trends that we observed. The Pujehun district had 49 confirmed EVD  
39  
40 464 cases. This number is much lower than in other districts. If it is true that the fear of Ebola may have  
41  
42 465 prevented people from accessing health services, the small number of EVD cases in the community  
43  
44 466 may have also raised confidence, leading to the increase of utilization rates after the initial drop. The  
45  
46 467 strength of this study is that it uses data from a remote rural district in Sierra Leone, with a 6-year  
47  
48 468 observational period. The pre, intra, and post-Ebola periods data, allowed a comparison between  
49  
50 469 trends. DWA was working in this community before the outbreak began, which gave an advantage of  
51  
52 470 knowledge of the setting when the epidemic began, which in turn facilitated mitigating measures to  
53  
54 471 be put in place. In addition, this allowed a collection of data in a prospective way, reducing the  
55  
56 472 potential bias in the accuracy of the data reported by other studies.<sup>6 14 32 35</sup>

57  
58 473 Failures in providing effective health care are associated with a chiefly vertical focus on outbreak  
59  
60 474 control.<sup>19-21</sup> The approach implemented in the Pujehun district worked on strengthening all the  
61  
62 475 components of the health system - governance, human resources, community involvement - before,  
63  
64 476 during and, after the epidemic.

The strengthening of the health system in the district, compared to other districts, allowed the containment of the epidemic and, above all, to maintain and strengthen MCH services as shown by the data reported in the paper. Health facilities in the district, both at community and hospital level, were able to maintain their services during the epidemic, overcoming public fear of Ebola and lack of confidence in service providers, which led to the public staying away from facilities in other districts in Sierra Leone.<sup>14</sup> In post-crisis situations, "windows of opportunity" are opened for redirecting the policies of the national health systems, renovating specific sectors (e.g. human resources, epidemiological surveillance systems, financing, etc.) and renewing services/practices at the operational level.<sup>48</sup> In Pujehun the implementation of an RS immediately after the acute Ebola phase reduced delays in patients accessing care and enabled a significant improvement in all MCH indicators at hospital level. Other studies have also found that using this window of opportunity to introduce systems such as performance based financing can also produce positive outcomes.<sup>49</sup> As Sierra Leone continues its recovery, there is a need to quantify the impact of the outbreak on MCH care to guide long-term strategies for MHC services. This study provides evidence on strategies to increase the resilience of fragile healthcare services and the importance of NGOs and government collaboration to bring about change.

**Figure 1** Study area, the Pujehun district in Sierra Leone.

**Figure 2** Maternal and pediatric admissions at hospital level.

**Figure 3** C-sections, deliveries, MDOCs, pediatric and maternal deaths at hospital level.

**Figure 4** ANC 1, ANC 4, deliveries, and family planning at community level.

1  
2  
3 505  
4  
5 506  
6  
7 507  
8  
9 508  
10  
11 509  
12 510  
13  
14 511  
15  
16 512  
17  
18 513  
19  
20 514  
21  
22 515  
23 516  
24  
25 517  
26  
27 518  
28  
29 519  
30  
31 520  
32  
33 521  
34 522  
35  
36 523  
37  
38 524  
39  
40 525  
41  
42 526  
43  
44 527  
45 528  
46  
47 529  
48  
49 530  
50  
51 531  
52  
53 532  
54 533  
55  
56 534  
57  
58 535  
59  
60 536  
537

**Acknowledgements** We are deeply grateful to all the staff of Pujehun district hospital and Peripheral Health Units, District Health Management, and the personnel of Doctors with Africa CUAMM who worked in Pujehun during the epidemic. We thank James Dean for his help with proof-reading.

**Contributors** GLQ, FT, and GP contributed to study design, literature review, data analysis, data interpretation, writing, and review of the final manuscript. LF, FDG, DP, and CM performed data analysis, data interpretation, and drafting of the manuscript. DB, SS, AK, BZ, and VP contributed to data collection and data interpretation. BSC, AS, WM, and SJ contributed to data interpretation and review of the final manuscript. All co-authors contributed to the improvement of the article.

**Funding** This work was undertaken and funded by Doctors with Africa CUAMM.

**Competing Interests** None declared.

**Patient consent for publication** Obtained.

**Ethics approval** Sierra Leone Ethics and Scientific Review Committee, Directorate of Policy, Planning and Information, Ministry of Health and Sanitation, Sierra Leone.

**Data sharing statement** All data underlying the findings described in the manuscript are fully available without restriction.

**Disclaimer** The views expressed in this publication are the sole responsibility of the authors and do not necessarily reflect the views of the affiliated organisations.

## REFERENCES

1. World Health Organization. *Ebola response roadmap. Situation reports*. WHO, 2018. Available from: <http://www.who.int/csr/disease/ebola/situation-reports/archive/en/> [Accessed March 2019].
2. Jones S, Ameh C. Exploring the impact of the Ebola outbreak on routine maternal health services in Sierra Leone, 2015. Available from: [https://www.vsointernational.org/sites/vso\\_international/files/vso\\_sierra\\_leone\\_\\_](https://www.vsointernational.org/sites/vso_international/files/vso_sierra_leone__) [Accessed March 2019].
3. Streifel C. *How did Ebola impact maternal and child health in Liberia and Sierra Leone? A report of the CSIS Global Health Policy Center*. CSIS, 2015.
4. Brolin Ribacke KJ, Saulnier DD, Eriksson A, *et al*. Effects of the West Africa Ebola Virus Disease on Health-Care Utilization - A Systematic Review. *Front Public Health* 2016;4:222.
5. Government of Sierra Leone, Ministry of Health and Sanitation. *Ebola virus disease situation report*. MoHS, 2015.
6. Brolin Ribacke KJ, van Duinen AJ, Nordenstedt H, *et al*. The Impact of the West Africa Ebola Outbreak on Obstetric Health Care in Sierra Leone. *PLoS One* 2016;11:e0150080.
7. Cancedda C, Davis SM, Dierberg KL, *et al*. Strengthening Health Systems While Responding to a Health Crisis: Lessons Learned by a Nongovernmental Organization During the Ebola Virus Disease Epidemic in Sierra Leone. *J Infect Dis* 2016; 214 (Suppl. 3):S153-S163.
8. Jones S, Sam B, Bull F, Pieh SB, *et al*. 'Even when you are afraid, you stay': Provision of maternity care during the Ebola virus epidemic: A qualitative study. *Midwifery* 2017;52:19-26.
9. Dynes MM, Miller L, Sam T, *et al*. Perceptions of the risk for Ebola and health facility use among health workers and pregnant and lactating women Kenema District, Sierra Leone, 2014. *Morb Mortal Wkly Rep* 2015;63:12267.
10. Delamou A, Hammonds RM, Caluwaerts S, *et al*. Ebola in Africa: beyond epidemics, reproductive health in crisis. *Lancet* 2014; 13; 384:2105.
11. Hayden EC. Ebola obstructs malaria control. *Nature* 2014;514:15-6.
12. Sierra Leone, Ministry of Health and Sanitation, United Nations Children's Fund. *Health facility survey. Assessing the impact of the EVD outbreak on health systems in Sierra Leone*. MoHS, UNICEF, 2014.
13. Sierra Leone, Ministry of Health and Sanitation, UN Children's Fund. *Sierra Leone health facility assessment 2015: impact of the EVD [Ebola Virus Disease] outbreak on Sierra Leone's primary health care system*. MoHS, UNICEF, 2015.

1  
2  
3 585 14. Jones S, Gopalakrishnan S, Ameh CA, *et al.* 'Women and babies are dying but not of Ebola': the  
4 586 effect of the Ebola virus epidemic on the availability, uptake and outcomes of maternal and  
5 587 newborn health services in Sierra Leone. *BMJ Glob Health* 2016;1(3):e000065.  
7 588  
8 589 15. Centers for Disease Control and Prevention (CDC). 2014-2016 Ebola Outbreak Distribution in West  
9 590 Africa. 2017. Available from:[https://www.cdc.gov/vhf/ebola/history/2014-2016-](https://www.cdc.gov/vhf/ebola/history/2014-2016-outbreak/distribution-map.html)  
10 591 [outbreak/distribution-map.html](https://www.cdc.gov/vhf/ebola/history/2014-2016-outbreak/distribution-map.html)[Accessed March 2019].  
12 592  
13 593 16. Sierra Leone, Ministry of Health and Sanitation. *Nationwide needs assessment for emergency*  
14 594 *obstetric and new-born care services in Sierra Leone*. MoHS, 2008.  
15 595  
16 596 17. Ajelli M, Parlamento S, Bome D, *et al.* The 2014 Ebola virus disease outbreak in Pujehun, Sierra  
18 597 Leone: epidemiology and impact of interventions. *BMC Med* 2015;13:281.  
19 598  
20 599 18. Quaglio GL, Pizzol D, Bome D, *et al.* Maintaining maternal and child health services during the Ebola  
21 600 outbreak: experience from Pujehun, Sierra Leone. *PLoS Currents* 2016;8:  
22 601 [ecurrents.outbreaks.d67aea257f572201f835772d7f188ba5](https://doi.org/10.1371/journal.ploscurrents.outbreaks.d67aea257f572201f835772d7f188ba5).  
24 602  
25 603 19. Dubois M, Wake C, Sturridge S, Bennett C. *The Ebola response in West Africa. Exposing the politics*  
26 604 *and culture of international aid*. 2015. Available from:  
27 605 <https://www.odi.org/sites/odi.org.uk/files/odi-assets/publications-opinion-files/9903.pdf>  
28 606 [Accessed March 2019]  
30 607  
31 608 20. Scott V, Crawford-Browne S, Sanders D. Critiquing the response to the Ebola epidemic through a  
32 609 primary health care approach. *BMC Public Health* 2016; 16:410 DOI 10.1186/s12889-016-3071-4.  
33 610  
34 611 21. Kruk ME, Myers M, Varpilah ST, *et al.* What is a resilient health system? Lessons from Ebola. *Lancet*  
35 612 2015;385:1910-12.  
36 613  
37 614 22. The Guardian. *Sierra Leone declares first Ebola free district*. 10 January 2015. Available  
38 615 from:<http://www.theguardian.com/world/2015/jan/10/sierraleonefirstebolafreedistrictwho>  
39 616 [Accessed March 2019].  
40 617  
41 618 23. Theuring S, Koroma AP, Harms G. "In the hospital, there will be nobody to pamper me": a  
42 619 qualitative assessment on barriers to facility-based delivery in post-Ebola Sierra Leone. *Reprod*  
43 620 *Health* 2018;15:155: 10.1186/s12978-018-0601-9.  
44 621  
45 622 24. Ly J, Sathananthan V, Griffiths T, *et al.* Facility-based delivery during the Ebola virus disease  
46 623 epidemic in rural Liberia: analysis from a cross-sectional, population-based household survey. *PLoS*  
47 624 *Med* 2016;13: e1002096.  
48 625  
49 626 25. Barden-O'Fallon J, Barry MA, Brodish P, *et al.* Rapid assessment of Ebola-related implications for  
50 627 reproductive, maternal, newborn, and child health service delivery and utilization in Guinea. *PLoS*  
51 628 *Curr* 2015;7:ecurrents.outbreaks.0b0ba06009d d091bc39ddb3c6d7b0826.  
52 629  
53 630 26. Lori JR, Rominski SD, Perosky JE, *et al.* A case series study on the effect of Ebola on facility-based  
54 631 deliveries in rural Liberia. *BMC Pregnancy Childbirth* 2015;15:254.  
55 632  
56 633  
57 634  
58 635  
59 636  
60 637


27. Iyengar P, Kerber K, Howe CJ, *et al.* Services for mothers and newborns during the Ebola outbreak in Liberia: the need for improvement in emergencies. *PLoS Curr* 2015; 7:ecurrents.outbreaks.4ba318308719ac86fbef91f8e56cb66f.
28. World Health Organization. *Situation report: Ebola virus disease, 10 June 2016*. WHO, 2018. Available from: [http://apps.who.int/iris/bitstream/10665/208883/1/ebolasitrep\\_10Jun2016\\_eng.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/208883/1/ebolasitrep_10Jun2016_eng.pdf?ua=1) [Accessed March 2019]
29. World Health Organisation. Monitoring emergency obstetric care. 2009 Available from: [https://apps.who.int/iris/bitstream/handle/10665/44121/9789241547734\\_eng.pdf;jsessionid=81643CFAAD44ECAEE6445C3BCBA40026?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/44121/9789241547734_eng.pdf;jsessionid=81643CFAAD44ECAEE6445C3BCBA40026?sequence=1) [Accessed March 2019]
30. R Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. Vienna, Austria, 2018.
31. Camara BS, Delamou A, Diro E, *et al.* Effect of the 2014/2015 Ebola outbreak on reproductive health services in a rural district of Guinea: an ecological study. *Trans R Soc Trop Med Hyg* 2017;11:22-9.
32. Delamou A, Hammonds RM, Caluwaerts S, *et al.* Effect of Ebola virus disease on maternal and child health services in Guinea: a retrospective observational cohort study. *Lancet Glob Health* 2017;5:e448-e457.
33. Statistics Sierra Leone, United Nations Children's Fund. *Sierra Leone multiple indicator cluster survey 2017*. Survey findings report. Stats SL, UNICEF, 2018.
34. Sierra Leone, Ministry of Health and Sanitation. *Sierra Leone demographic and health survey*. MoHS, 2013.
35. Wagenaar BH, Augusto O, Beste J, *et al.* The 2014-2015 Ebola virus disease outbreak and primary healthcare delivery in Liberia: Time-series analyses for 2010-2016. *PLoS Med* 2018;15:e1002508.
36. Nobles J, Frankenberg E, Thomas D. The Effects of mortality on fertility: population dynamics after a natural disaster. *Demography* 2015;52:15-38.
37. Hapsari ED, Widyawati, Nisman WA, *et al.* Change in contraceptive methods following the Yogyakarta earthquake and its association with the prevalence of unplanned pregnancy. *Contraception* 2009;79:316-22.
38. McBain RK, Wickett E, Mugunga JC, *et al.* The post-Ebola baby boom: time to strengthen health systems *Lancet* 2016;388: 2331-3.
39. Bolkan HA, Bash-Taqi DA, Samai M, *et al.* Ebola and indirect effects on health service function in Sierra Leone. *PLoS Curr* 2014;6:ecurrents.outbreaks.0307d588d f619f9c9447f8ead5b72b2d.



1  
2  
3 678 40. Olu O, Kargbo B, Kamara S, *et al.* Epidemiology of Ebola virus disease transmission among health  
4 679 care workers in Sierra Leone, May to December 2014: a retrospective descriptive study. *BMC Infect*  
5 680 *Dis* 2015;15:416.  
6 681  
7 682 41. Thaddeus S, Maine D. Too far to walk: maternal mortality in context. *Sm Sci Med* 1994;38:1091-  
8 683 1110. 41.  
9 684  
10 685 42. Treacy L. Distance, accessibility and costs. Decision-making during childbirth in rural Sierra Leone:  
11 686 A qualitative study. *PLoS One* 2018;13: e0188280.  
12 687  
13 688 43. Tayler-Smith K, Zachariah R, Manzi M, *et al.* An ambulance referral network improves access to  
14 689 emergency obstetric and neonatal care in a district of rural Burundi with high maternal mortality.  
15 690 *Trop Med Int Health* 2013;18:993-1001.  
16 691  
17 692 44. Groppi L, Somigliana E, Pisani V, *et al.* A hospital-centred approach to improve emergency obstetric  
18 693 care in South Sudan. *Int J Gynaecol Obstet* 2015;128:58–61.  
19 694  
20 695 45. Tsegaye A, Somigliana E, Alemayehu T, *et al.* Ambulance referral for emergency obstetric care in  
21 696 remote settings. *Int J Gynaecol Obstet* 2016;133:316-9.  
22 697  
23 698 46. Oyerinde K, Harding Y, Amara P, *et al.* Barriers to uptake of emergency obstetric and newborn care  
24 699 services in Sierra Leone: a qualitative study. *Comm Med & Health Educ* 2012;2: 10.4172/2161-  
25 700 0711.1000149.  
26 701  
27 702 47. Vallières F, Cassidy EL, McAuliffe E, *et al.* Can Sierra Leone maintain the equitable delivery of their  
28 703 Free Health Care Initiative? The case for more contextualised interventions: results of a cross-  
29 704 sectional survey. *BMC Health Serv Res* 2016;16:258.  
30 705  
31 706 48. Witter S, Hunter B. *Resilience of health systems during and after crises – what does it mean and*  
32 707 *how can it be enhanced?* ReBUILD Research Programme Consortium. 2017. Available  
33 708 from:[https://rebuildconsortium.com/media/1535/rebuild\\_briefing\\_1\\_june\\_17\\_resilience.pdf](https://rebuildconsortium.com/media/1535/rebuild_briefing_1_june_17_resilience.pdf)  
34 709 [Accessed March 2019].  
35 710  
36 711 49. Mussah VG, Mapleh L, Ade S, *et al.* Performance-based financing contributes to the resilience of  
37 712 health services affected by the Liberian Ebola outbreak. *Public Health Action* 2017;7(Suppl.  
38 713 1):S100-S105.  
39 714  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60









1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

- C-sections
- Deliveries
- Maternal deaths
- MDOCs
- Pediatric deaths





## Maternal admissions, maternal deaths, C-sections, and MDOCs at hospital level.

	Maternal admissions			Maternal deaths			C-sections			MDOC		
	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value
Difference between average of Ebola period vs pre-Ebola period	7	-7 to 22	0.333	-1	-2 to 0	0.042	5	-1 to 11	0.13	2	-11 to 14	0.782
Difference between average of post-Ebola period vs Ebola period	43	28 to 58	<0.001	2	1 to 3	0.001	15	8 to 21	<0.001	41	30 to 54	<0.001
Difference between average of post-Ebola period vs pre-Ebola period	50	37 to 64	<0.001	1	0 to 2	0.135	19	13 to 25	<0.001	43	31 to 54	<0.001
<b>Pre-Ebola period</b>												
Number of events over pre-Ebola period ( $\beta_0$ )	49	37 to 61	<0.001	1	0 to 2	0.026	9	4 to 14	0.001	16	5 to 26	0.003
Trend in number over pre-Ebola period ( $\beta_1$ )	0	0 to 1	0.281	0	0 to 0	0.677	0	-0 to 0	0.999	0	0 to 0.5	0.768
<b>Ebola period</b>												
Average monthly change in number over Ebola period ( $\beta_2$ )	-40	-60 to -19	<0.001	0	-2 to 0	0.480	2	-7 to 11	0.668	-11	-29 to 6	0.207
Difference between trend of Ebola period vs pre-Ebola period ( $\beta_3$ )	7	4 to 11	<0.001	0	0 to 0	0.605	1	-1 to 2	0.346	4	1 to 7	0.006
<b>Post-Ebola period</b>												
Average monthly change in number during post-Ebola period ( $\beta_4$ )	11	-7 to 30	0.23	1	0 to 2	0.258	13	5 to 21	0.001	16	0 to 32	0.044
Difference between trend of post-Ebola period vs Ebola period ( $\beta_5$ )	-7	-10 to -4	<0.001	0	0 to 0	0.665	-1	-2 to 0.8	0.433	-4	-7 to -1	0.009
Difference between trend of post-Ebola vs pre-Ebola period ( $\beta_3 + \beta_5$ )	0	-1 to 1	1	0	0 to 0	0.657	0	0 to 0	0.431	0	0 to 1	0.503

## Pediatric admissions, pediatric deaths, and institutional deliveries at hospital level.

	Pediatric admissions			Pediatric deaths			Institutional deliveries		
	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value
Difference between average of Ebola period vs pre-Ebola period	1	-39 to 40	0.968	-1	-6 to 5	0.826	11	2 to 21	0.02
Difference between average of post-Ebola period vs Ebola period	133	92 to 174	<0.001	9	3 to 15	0.004	28	18 to 38	<0.001
Difference between average of post-Ebola period vs pre-Ebola period	134	98 to 170	<0.001	8	3 to 14	0.003	39	31 to 48	<0.001
<b>Pre-Ebola period</b>									
Number of events over pre-Ebola period ( $\beta_0$ )	46	10 to 82	0.011	7	2 to 12	0.007	27	19 to 34	<0.001
Trend in number over pre-Ebola period ( $\beta_1$ )	0	-2 to 2	0.808	0	0 to 0	0.641	0	0 to 0	0.42
<b>Ebola period</b>									
Average monthly change in number over Ebola period ( $\beta_2$ )	1	-48 to 50	0.955	1	-7 to 9	0.836	-12	-25 to 1	0.072
Difference between trend of Ebola period vs pre-Ebola period ( $\beta_3$ )	1	-8 to 10	0.823	0	-1 to 2	0.763	4	2 to 6	0.001
<b>Post Ebola period</b>									
Average monthly change in number over post-Ebola period ( $\beta_4$ )	53	5 to 100	0.029	6	-1 to 14	0.086	11	-1 to 22	0.064
Difference between trend of post-Ebola period vs Ebola period ( $\beta_5$ )	2	-7 to 10	0.702	0	-1 to 1	0.899	-3	-5 to -1	0.001
Difference between trend of post-Ebola vs pre-Ebola period ( $\beta_3 + \beta_5$ )	3	0 to 5	0.035	0	0 to 0	0.423	0	0 to 0	0.486

## Institutional delivery, ANC 1, ANC 4 and family planning at community level.

	Institutional delivery			ANC 1			ANC 4			Family planning		
	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value
Difference between average of Ebola period vs pre-Ebola period	148	99 to 196	<0.001	74	3 to 145	0.042	80	21 to 139	0.008	490	-92 to 1073	0.099
Difference between average of post-Ebola period vs Ebola period	-10	-59 to 39	0.695	-48	-122 to 26	0.2	23	-38 to 84	0.461	-262	-855 to 330	0.386
Difference between average of post-Ebola period vs pre-Ebola period	138	93 to 183	<0.001	26	-40 to 91	0.448	103	48 to 157	<0.001	228	-293 to 750	0.391
<b>Pre Ebola period</b>												
Number of events over pre-Ebola period ( $\beta_0$ )	688	643 to 732	<0.001	1062	1002 to 1121	<0.001	694	644 to 743	<0.001	2690	2187 to 3193	<0.001
Trend in number over pre-Ebola period ( $\beta_1$ )	8	6 to 10	<0.001	7	4 to 10	<0.001	6	4 to 8	<0.001	69	42 to 95	<0.001
<b>Ebola period</b>												
Average monthly change in number over Ebola period ( $\beta_2$ )	-28	-90 to 34	0.382	-61	-161 to 40	0.238	-94	-176 to -11	0.027	-671	-1431 to 89	0.084
Difference between trend of Ebola period vs pre-Ebola period ( $\beta_3$ )	-1	-12 to 10	0.881	-5	-21 to 12	0.591	5	-8 to 19	0.437	-26	-156 to 104	0.692
<b>Post Ebola period</b>												
Average monthly change in number during post-Ebola period ( $\beta_4$ )	-25	-81 to 30	0.37	-5	-94 to 83	0.906	35	-37 to 109	0.343	-51	-759 to 657	0.888
Difference between trend of post-Ebola period vs Ebola period ( $\beta_5$ )	-7	-17 to 4	0.228	-2	-18 to 15	0.819	-13	-27 to 0	0.056	-59	-186 to 68	0.361
Difference between trend of post-Ebola vs pre-Ebola period ( $\beta_3 + \beta_5$ )	-7	-10 to -4	<0.001	-6	-10 to -3	<0.001	-8	-11 to -5	<0.001	-85	-119 to -51	<0.001

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No.	Recommendation	Page No.	Relevant text from manuscript
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	1	
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2	
Introduction				
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4-6	
Objectives	3	State specific objectives, including any prespecified hypotheses	6	
Methods				
Study design	4	Present key elements of study design early in the paper	7	
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	7-8	
Participants	6	(a) <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	8	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6	
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	7-8	
Bias	9	Describe any efforts to address potential sources of bias	16-17	
Study size	10	Explain how the study size was arrived at		

Continued on next page

Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	9-10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	10
		(b) Describe any methods used to examine subgroups and interactions	10
		(c) Explain how missing data were addressed	NA
		(d) <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy NA	NA
		(e) Describe any sensitivity analyses	NA
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	9-12
		(b) Give reasons for non-participation at each stage	NA
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	7
		(b) Indicate number of participants with missing data for each variable of interest	
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	9-11
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	9-11
		(b) Report category boundaries when continuous variables were categorized	9-11
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	9-11

Continued on next page

Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	NA
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	13
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	16-17
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	16-17
Generalisability	21	Discuss the generalisability (external validity) of the study results	16-17
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	18

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

# BMJ Open

## Impact of Ebola outbreak on reproductive health services in a rural district of Sierra Leone. A prospective observational study.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-029093.R2
Article Type:	Research
Date Submitted by the Author:	15-May-2019
Complete List of Authors:	<p>Quaglio , GianLuca ; European Parliament  Tognon, Francesca; Department for Woman and Child Health, University of Padua, Italy  Finos, Livio; Department of Developmental Psychology and Socialisation, University of Padua, Italy  Bome, David; Ministry of Health and Sanitation, Freetown, Sierra Leone  Sesay, Santigie ; Ministry of Health and Sanitation, Freetown, Sierra Leone  Atiba, Kebbie ; Department for Woman and Child Health, Pujehun Hospital, Sierra Leone  Di Gennaro, Francesco ; Department of Infectious Diseases, University of Bari, Italy  Bienvenu Salim, Camara ; National Centre for Training and Research in Rural Health of Maferinyah, Forécariah, Guinea  Marotta, Claudia; Department of Science for Health Promotion and Mother to Child Care, University of Palermo, Italy  Pisani, Vincenzo; Department for Woman and Child Health, Pujehun Hospital, Sierra Leone, Pujehun, SL  Bangura , Zainab; Department for Woman and Child Health, Pujehun Hospital, Sierra Leone  Pizzol, Damiano; Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy  Saracino, Annalisa; Department of Infectious Diseases, University of Bari, Italy  MAZZUCCO, WALTER; Department of Science for Health Promotion and Mother to Child Care, University of Palermo, Italy  Jones, Susan; Department of Nursing and Midwifery School of Human and Health Sciences, University of Huddersfield, UK  Putoto, G; Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy</p>
<b>Primary Subject Heading</b>:	Health policy
Secondary Subject Heading:	Public health, Mental health, Infectious diseases, Addiction, Communication
Keywords:	PAEDIATRICS, Public health < INFECTIOUS DISEASES, Community gynaecology < GYNAECOLOGY



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



SCHOLARONE™  
Manuscripts

# Impact of Ebola outbreak on reproductive health services in a rural district of Sierra Leone. A prospective observational study.

Gianluca Quaglio,<sup>1,2,3</sup> Francesca Tognon,<sup>4</sup> Livio Finos,<sup>5</sup> David Bome,<sup>6</sup> Sesay Santigie,<sup>6</sup> Atiba Kebbie,<sup>7</sup> Francesco Di Gennaro,<sup>8</sup> Bienvenu Salim Camara,<sup>9</sup> Claudia Marotta,<sup>10</sup> Vincenzo Pisani,<sup>7</sup> Bangura Zainab,<sup>7</sup> Damiano Pizzol,<sup>3</sup> Annalisa Saracino,<sup>8</sup> Walter Mazzucco,<sup>10</sup> Susan Jones,<sup>11</sup> Giovanni Putoto<sup>3</sup>

1. European Parliamentary Research Services (EPRS), European Parliament, Brussels, Belgium;
2. Department of International Health/CAPHRI, Maastricht University, The Netherlands;
3. Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy;
4. Department for Woman and Child Health, University of Padua, Italy;
5. Department of Developmental Psychology and Socialisation, University of Padua, Italy;
6. Ministry of Health and Sanitation, Freetown, Sierra Leone;
7. Department for Woman and Child Health, Pujehun Hospital, Sierra Leone;
8. Department of Infectious Diseases, University of Bari, Italy;
9. National Centre for Training and Research in Rural Health of Maferinyah, Forécariah, Guinea;
10. Department of Science for Health Promotion and Mother to Child Care, University of Palermo, Italy;
11. Department of Nursing and Midwifery School of Human and Health Sciences, University of Huddersfield, UK.

## Corresponding author

Gianluca Quaglio, MD  
Directorate-General for Parliamentary Research Services (EPRS)  
European Parliament  
Rue Wiertz, 60, B-1047  
Brussels, Belgium  
gianluca.quaglio@europarl.europa.eu

**Words count** (excluding title page, abstract, references, figures and tables): 5182



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**ABSTRACT**

**Objectives** To assess the trends concerning utilisation of maternal and child health (MCH) services before, during, and after the Ebola outbreak, quantifying the contribution of a reorganised referral system (RS).

**Design** A prospective observational study of MCH services.

**Setting** Pujehun district in Sierra Leone, 77 community health facilities and 1 hospital from 2012 to 2017.

**Main outcome measures** MCH utilization was evaluated by assessing: i) institutional deliveries, Cesarean-sections, paediatric and maternity admissions and deaths, and major direct obstetric complications (MDOCs), at hospital level; ii) antenatal care (ANC) 1 and 4, institutional delivery, and family planning, at community level. Contribution of a strengthened RS was also measured.

**Results** At hospital level, there is a significant difference between trends Ebola vs pre-Ebola for maternal admissions (7, 95% CI 4 to 11,  $p < 0.001$ ), MDOCs (4, 95% CI 1 to 7,  $p = 0.006$ ), and institutional deliveries (4, 95% CI 2 to 6,  $p = 0.001$ ). There is also a negative trend in the transition from Ebola to post Ebola for maternal admissions (-7, 95% CI -10 to -4,  $p < 0.001$ ), MDOCs (-4, 95% CI -7 to -1,  $p 0.009$ ) and institutional deliveries (-3, 95% CI -5 to -1,  $p 0.001$ ). The differences between trends pre-Ebola vs post-Ebola are only significant for pediatric admissions (3, 95% CI 0 to 5,  $p 0.035$ ). At community level, the difference between trends Ebola vs pre-Ebola and Ebola vs post-Ebola are not significant for any indicators. The differences between trends pre-Ebola vs post-Ebola show a negative difference for institutional deliveries (-7, 95% CI -10 to -4,  $p < 0.001$ ) ANC 1 (-6, 95% CI -10 to -3,  $p < 0.001$ ), ANC 4 (-8, 95% CI -11 to -5,  $p < 0.001$ ) and family planning (-85, 95% CI -119 to -51,  $p < 0.001$ ).

**Conclusions** A stronger health system compared to other districts in Sierra Leone and a strengthened RS enabled health facilities in Pujehun to maintain service provision and uptake during and after the Ebola epidemic.

**Keywords:** Ebola, Sierra Leone, Maternal and Child Health indicators, Referral system, Reproductive health service.

## Strengths and limitations of this study

- ▶ The study uses data from a remote rural district in Sierra Leone, with a 6-year observational period. Data have been collected in a prospective way, reducing the potential bias in the accuracy of the data reported by other studies carried out in countries affected by Ebola.
- ▶ Data from pre, intra, and post-Ebola periods allowed comparisons between trends, something rarely carried out in countries heavily affected by Ebola.
- ▶ The data refers to a single area of Sierra Leone: the sample cannot be considered representative of the country as a whole.
- ▶ In addition to measures put in place to reduce the impact of the disease on mothers and children, Pujehun had far fewer Ebola cases than other districts, which may also have led to the utilization of health services.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**INTRODUCTION**

The 2014-2015 Ebola Virus Disease (EVD) outbreak was the most severe in history, mainly affecting three West African countries; Guinea, Sierra Leone and Liberia. Overall 28,616 people were infected of which 11,310 died and the outbreak was declared a global public health emergency by the WHO.<sup>1</sup> Of the three countries affected, Sierra Leone had the most confirmed cases (8,704), which accounted for 50% of all confirmed cases in West Africa, and 3,589 deaths.<sup>2-4</sup> All 14 districts in Sierra Leone were affected, but at different times and to varying degrees.<sup>5</sup> During the Ebola crisis the population's trust in the national health system declined in Sierra Leone, leading to an overall reduction in the use of health services, including reproductive, maternal, and child services.<sup>6-8</sup> Underlying factors for the decrease in the use of health services included fear of infection, for both healthcare workers and patients, the underlying fragility of the health systems, the reduced numbers of available health personnel, and the death of healthcare workers due to EVD.<sup>9 10</sup> It has been estimated that 30% of health workers who died of EVD in West Africa were maternal and child healthcare (MCH) providers.<sup>11</sup> However, there were considerable variations in the reduction of health service uptake when looked at by district level in Sierra Leone.<sup>6 12-14</sup> While districts such as Kambia, Port Loko and Bonthe showed large reductions in facility-based delivery (between 38-41%), the district of Pujehun showed only a 5% decrease in the same service. Similar geographic variations were seen in the reduction in antenatal care (ANC) visits.<sup>12 13</sup>

The number of confirmed EVD cases - and deaths - varied considerably by district. There were no more than 100 confirmed cases in both Bonthe and Pujehun, and up to 4,000 confirmed cases in both Port Loko and Bombali.<sup>15</sup> However, public fear of Ebola, regardless of the actual number of cases per district, may still have prevented many people from accessing services. The challenge of providing adequate levels of care during a humanitarian emergency such as the EVD crisis was further exacerbated by the weak health system in Sierra Leone, particularly in rural areas where the poor condition of the roads and high transport costs cause delays in accessing services, and contribute to increased maternal and neonatal mortality.<sup>16</sup>

Doctors with Africa (DwA) CUAMM is an Italian NGO working in Sierra Leone since 2012. It is present in the Pujehun district focusing on MCH care both at hospital and community level<sup>17 18</sup> In this paper, community level refers to Peripheral Health Units (PHUs), i.e. all health facilities outside the hospital. As described in our previous reports,<sup>17 18</sup> a number of measures were put in place to control the Ebola epidemic in the Pujehun district which might have reduced the impact of the disease on mothers and children compared to other districts. During this EVD epidemic, the predominantly

vertical focus on outbreak control was associated with failures in providing effective care for routine health needs.<sup>19-21</sup> In contrast, the approach implemented in the Pujehun district was not based on vertical actions and 'humanitarian response to health emergencies with a short half-life'.<sup>21</sup> Rather, it worked on strengthening all the components of the health system - governance, human resources, community involvement - before, during and, after the epidemic. A rapid response to the crisis by the local health authorities was implemented adopting public health measures before any other district in Sierra Leone.<sup>22</sup> The activities were mainly concentrated on keeping the health service open and properly functioning in order to reduce the collateral effects of the epidemic on routine health services. No health units in the Pujehun district were closed during the epidemic. Measures to empower community leaders and use culturally appropriate methods of communication helped to dispel community mistrust in the health services. At community level, a number of strategies were implemented such as the regular rotation of health facility staff, which strengthened teamwork and effective leadership. In Sierra Leone, healthcare workers based at community health centres may often work alone in isolated centres with limited support from clinical colleagues or management. By rotating staff through the various facilities, they gain on the job training, peer support, and develop new working relationships. At the start of the Ebola epidemic, many expatriate healthcare workers in NGOs left Sierra Leone, negatively affecting care delivery and staff morale. The continued presence of international teams in the daily activities in Pujehun hospital and the acceptance of the professional risks by both national and international staff may have contributed to maintaining an attitude of 'normality' in an extremely stressful environment. This might also help to explain the population's positive receptiveness towards the health services.<sup>17 18</sup>

Different types of referral systems (RSs) such as motorbikes were present in the country in the pre Ebola period to transport patients from the villages to the nearest health facility. Ambulances were also present in several districts with 73% of health facilities nationwide having a functioning RS, 59% of them consisting of an ambulance on call.<sup>12 23</sup> In the Pujehun district, the RS was barely functioning, only able to support the activity of a limited number of PHUs. The service was also entirely funded by the patients themselves, resulting in underutilization of the service. Utilization was further reduced during the outbreak, when the ambulances were identified by the population with the transport of Ebola infected patients, and their use occasioned fear and distrust. In January 2015, in collaboration with the Ministry of Health and Sanitation (MoHS) of Sierra Leone and UNICEF, DWA began the re-organisation and reinforcement of the RS, transferring pregnant women and pediatric cases from PHUs to the Pujehun hospital.

Our previous studies<sup>18</sup> provided information only on three MCH indicators, namely pediatric admissions, maternity admissions, and institutional deliveries; in addition it did not assess the trends in the post-EVD period. Existing studies examining the influence of EVD on MCH services targeted the outbreak and the immediate post-outbreak periods.<sup>24-27</sup> Understanding the trends in the use of MCH services before, during, and after the EVD outbreak will help to guide post-EVD interventions, increasing access to MCH services in rural Sierra Leone. This information will also be useful in preparing a more organised and structured RS. With this background, the aims of this study are: i) to assess trends in institutional deliveries, C-sections, paediatric and maternity admissions, paediatric and maternity deaths, and major direct obstetric complications (MDOCs), before, during, and after the EVD in the Pujehun hospital, thus complementing the results of the previous report which were limited to 3 MCH indicators; ii) to assess trends in ANC 1 and 4, institutional delivery, and family planning, at community level. This study was carried out in conjunction with the strengthening of an RS initiated a few weeks after the Pujehun district was declared Ebola-free.

METHODS

Setting

Sierra Leone has four provinces that are divided into 14 districts. Pujehun is one of four districts in the southern province (Figure 1). It has a population of approximately 375,000 inhabitants. The primary care network included 77 MoHS PHUs, 5 of which provide basic emergency obstetrics care (BEmOC). The secondary care system consists of the MoHS provided district hospital, which comprises the MCH complex, providing comprehensive emergency obstetric and newborn care (CEmONC) services. Connections between the community and health facilities are difficult because of the very poor condition of the roads. Furthermore, the district is divided by a major river (Moa River) and has a riverine area reachable only with boats, which further hinders access. The first case of Ebola in Pujehun district was reported on the 7th July 2014. The district was declared Ebola free on the 10th January 2015.<sup>28</sup> A total of 49 patients were registered with a case fatality rate of 85.7% (42/49).

Referral system

In the Pujehun district, two ambulances managed by the District Health Management Team (DHMT) were functioning in the pre Ebola period, but only 63% of the PHUs were able to use the service.<sup>12 23</sup> Emergency calls were not coordinated by the hospital and the transport costs were covered by the patients, dissuading many from using the service. During the outbreak, people came to associate the

ambulances with transporting Ebola infected patients, which further discouraged their use. A 24-h free-of-charge ambulance RS, transferring pregnant women with obstetric complications from the health centers to Pujehun hospital was implemented in January 2015. In the hospital a call center was established and the call center number was distributed to all the 77 PHUs. Private calls were considered only in the case of an emergency or if the staff of the PHU were not available. After confirming an emergency condition together with the PHU staff, the hospital midwife had the responsibility to authorize the referral. A nurse on duty from the maternity hospital accompanied the driver in each referral. Health personnel at hospital and PHUs levels were trained on Life Saving Skills – Emergency Obstetric and Newborn Care, including referral criteria and definition of MDOCs.<sup>29</sup>

Referrals were carried out by 3 ambulances, two positioned in the Pujehun MCH complex, and a third one in Jendema, bordering Liberia, on the opposite side of the Moa River. Around the Jendema area, 15 PHUs were located serving a population of approximately 80,000 inhabitants. Referrals in this area were made using the ambulances and by transferring patients at the river crossing point via a barge or a motor boat, depending on the flow rate of the river. Pediatric referrals were performed using private motor bikes available in the villages and hired from PHUs staff without the involvement of the call center. A referral form describing the clinical case and the justification for the referral was distributed to all the PHUs. The bike rider, after bringing the patient to the pediatric ward, delivered the referral form and received the reimbursement. For all patients carried to the hospital information was collected, including demographics, location, and the reason for contacting the RS. Community awareness activities were organized about the RS through meetings and radio discussions held by the DMHT, hospital health personnel, and local authorities.

### **Study design, population, and period**

A prospective observational study using routinely collected health services data, from January 2012 to December 2017, was carried out. Three time periods were considered: pre- Ebola period (1<sup>st</sup> January 2012 – 30<sup>th</sup> May 2014); Ebola period (1<sup>st</sup> June 2014 – 28<sup>th</sup> February 2015); post- Ebola period (1<sup>st</sup> March 2015 – 31<sup>th</sup> December 2017). We considered the Ebola period from one month before the first confirmed case in the district (i.e. June 2014), to one month after the country being declared Ebola free (i.e. February 2015). This was done because in Sierra Leone the outbreak had started in other districts of the country before the first case registered in Pujehun and continued to affect other districts until November 2015. It is realistic to assume that public fear of potential EVD cases and lack of confidence in the health services persisted in the Pujehun population during that time.<sup>14</sup> In



1  
2  
3 220 addition, expanding the Ebola period enabled a full assessment of the impact of the disease with an  
4  
5 221 adequate comparison with the two long periods before and after the Ebola epidemic.  
6  
7 222  
8  
9 223 **Data collection**  
10  
11 224 Data on MCH indicators was prospectively collected from hospital registers (maternity ward, delivery  
12  
13 225 unit, pediatric ward, operating theatre). The following variables were collected on a monthly basis:  
14  
15 226 1) paediatrics admissions; 2) pediatric deaths; 3) maternity admissions; 4) maternal deaths; 5)  
16  
17 227 deliveries; 6) C-sections; 7) MDOC cases. MDOC cases were collected using a dedicated database  
18  
19 228 within the hospital and confirmed by a gynaecologist. All hospital maternal deaths were reviewed by  
20  
21 229 DHMT and classified according to Maternal Death Surveillance and Response policy by MoHS.  
22  
23 230 Paediatric deaths did not include stillbirths and early neonatal deaths, but only deaths of children  
24  
25 231 admitted to the paediatric ward.  
26  
27 232 At community level, the following variables were collected from the local district Health  
28  
29 233 Management Information System (HMIS): 1) family planning consultations per month; 2) deliveries  
30  
31 234 per month; 3) ANC 1 per month; 4) ANC 4 per month. Different variables were collected from the  
32  
33 235 two types of sites, based on the different services provided at community level (BEmOC) and at  
34  
35 236 hospital level (CEmONC). Quarterly review meetings were organized with the staff in charge of the  
36  
37 237 health facilities to address data discrepancies in the reports. Technical assistance was provided to  
38  
39 238 the DHMT to improve timeliness, completeness, and accuracy of data regarding CEmOC and BEmONC  
40  
41 239 services.  
42  
43 240 For the RS, data was collected from records of all of the study sites, including delivery registers,  
44  
45 241 delivery logbooks, prenatal registers, referral registers, and death registers. Additional data was  
46  
47 242 collected from the ambulance database and logbook. Records in the database were then validated  
48  
49 243 by cross-checking the records with registers at the study sites.  
50  
51 244  
52  
53 245 **Statistical analysis**  
54  
55 246 For each indicator, a segmented seasonal autoregressive model of order 1 was estimated. The  
56  
57 247 segments defined the three periods: before the EVD epidemic (January 2012 to May 2014), during  
58  
59 248 the epidemic (June 2014 to February 2015), and after the epidemic (March 2015 to December 2017).  
60  
Differences were considered statistically significant at  $p < 0.05$ . The analysis was performed using  
R.<sup>30</sup> The full description of the methodology of the statistical analysis is available in Annex 1.

## Patient involvement

No patients were involved in defining the research question or the outcome measures, nor were they involved in the design and implementation of the study. There are no plans to involve patients in the dissemination of the results. The full statistical analysis is available in Annex 2.

## RESULTS

### Hospital level: Pre-Ebola period

At hospital level, for all indicators, the trend is stable during the pre Ebola period, without significant changes (Figure 2 and 3).

### Hospital level: Ebola vs pre-Ebola period

At hospital level, the differences between Ebola period vs pre-Ebola averages show a statistically significant increase for institutional deliveries (11, 95% CI 2 to 21,  $p = 0.02$ ) and for the reduction of maternal deaths (-1, 95% CI - 2 to 0,  $p = 0.042$ ) (Table 1). There is also a statistically significant difference between the trend of Ebola period vs pre-Ebola period, for maternal admissions (7, 95% CI 4 to 11,  $p < 0.001$ ), MDOCs (4, 95% CI 1 to 7,  $p = 0.006$ ), and institutional deliveries (4, 95% CI 2 to 6,  $p = 0.001$ ) (Figure 2 and 3).

### Hospital level: Ebola vs post-Ebola period

At hospital level, the differences between averages of the post Ebola vs Ebola are statistically significant for all indicators: institutional deliveries, C-sections, paediatric and maternity admissions, paediatric and maternity deaths, and MDOCs (Table 1). There is also a negative trend in the transition from Ebola to post Ebola for maternal admissions (-7, 95% CI -10 to -4,  $p < 0.001$ ), MDOCs (-4, 95% CI -7 to -1,  $p = 0.009$ ) and institutional deliveries (-3, 95% CI -5 to -1,  $p = 0.001$ ) (Figure 2 and 3).

### Hospital level: Pre-Ebola vs post-Ebola period

The differences between averages of the pre-Ebola vs post-Ebola periods are also statistically significant for all indicators, except for maternal deaths (Table 1). The differences between trends between pre-Ebola vs post-Ebola period are only significant for pediatric admissions (3, 95% CI 0 to 5,  $p = 0.035$ ) (Figure 2 and 3).

Table 1 MCH indicators at hospital and community level									
Indicator	Difference between average of Ebola period vs pre-Ebola period			Difference between average of Ebola period vs post-Ebola period			Difference between average of pre-Ebola period vs post-Ebola period		
	β	95% CI	p value	β	95% CI	p value	β	95% CI	p value
<b>HOSPITAL LEVEL</b>									
Maternal admissions	7	-7 to 22	0.333	43	28 to 58	<0.001	50	37 to 64	<0.001
Maternal deaths	-1	-2 to 0	0.042	2	1 to 3	0.001	1	0 to 2	0.135
Institutional deliveries	11	2 to 21	0.02	28	18 to 38	<0.001	39	31 to 48	<0.001
C-sections	5	-1 to 11	0.13	15	8 to 21	<0.001	19	13 to 25	<0.001
MDOC	2	-11 to 14	0.782	41	30 to 54	<0.001	43	31 to 54	<0.001
Pediatric admissions	1	-39 to 40	0.968	133	92 to 174	<0.001	134	98 - 170	<0.001
Pediatric deaths	-1	-6 to 5	0.826	9	3 to 15	0.004	8	3 to 14	0.003
<b>COMMUNITY LEVEL</b>									
Institutional deliveries	148	99 to 196	<0.001	-10	-59 to 39	0.695	138	93 to 183	<0.001
ANC 1	74	3 to 145	0.042	-48	-122 to 26	0.2	26	-40 to 91	0.448
ANC 4	80	21 to 139	0.008	23	-38 to 84	0.461	103	48 to 157	<0.001
Family planning	490	-92 to 1073	0.099	-262	-855 to 330	0.386	228	-293 to 750	0.391

Community level: Pre-Ebola period

At community level, all indicators in the months before Ebola showed a positive trend. There was a monthly average increase of 8 institutional deliveries (95% CI 6 to 10,  $p<0.001$ ); a monthly average increase of 7 ANC 1 (95% CI 4 to 10,  $p<0.001$ ) and 6 ANC 4 (95% CI 4 to 8,  $p<0.001$ ), and a monthly average increase of 69 women accessing family planning services (95% CI 42 to 95,  $p<0.001$ ) (Figure 4).

Community level: Ebola vs pre-Ebola period

At community level, with the exception of family planning, the differences between averages of Ebola period vs pre-Ebola are statistically significant for all indicators: institutional deliveries (148, 95% CI 99 to 196,  $p<0.001$ ), ANC 1 (74, 95 % CI 3 to 145,  $p=0.042$ ), and ANC 4 (80, 95% CI 21 to 139,  $p=0.008$ ) (Table 1). The difference between trends (Figure 3) of the Ebola vs pre-Ebola period are not significant for any of the indicators considered (Figure 4).

Community level: Ebola vs post-Ebola period

At community level, the differences between averages (Table 1) and the difference between trends (Figure 4) of the Ebola vs post-Ebola period are not significant for any of the indicators considered.

Community level: Pre-Ebola vs post-Ebola period

The differences between averages of the pre-Ebola vs post-Ebola are statistically significant, with an increase in institutional deliveries (138, 95% CI 93 to 183,  $p<0.001$ ) and ANC 4 (103, 95% CI 48 to 157,

p < 0.001) (Table 1). However, there is a negative difference between trends among the two periods, for all the variables considered: institutional deliveries (-7, 95% CI -10 to -4, p < 0.001) ANC 1 (-6, 95% CI -10 to -3, p < 0.001), ANC 4 (-8, 95% CI -11 to -5, p < 0.001) and most significantly for family planning (-85, 95% CI -119 to -51, p < 0.001) (Figure 4).

### **Referral system: Obstetric and paediatric results**

Between January 2015 and December 2017 there were 2,450 obstetric referrals. Of these, 1,574 (64%) were MDOC, which represent 70% of all the 2,233 MDOCs treated in the hospital over the same period. The baseline characteristics and reasons for MDOCs collected through the RS are reported on Table 2. At the same time, 4,671 paediatric patients were admitted in the hospital through the RS, representing 72% of the 6,518 total admission during the same period. Reasons for paediatric referrals are shown on Table 3.

Table 2 Baseline characteristics and reasons for MDOCs collected through RS, period 2015 - 2017			
Age (years)		N	%
Mean		25,3	SD 7
12-19		442	28%
20-29		613	39%
30-39		464	29%
40+		43	3%
Unknown		12	1%
Number of previous deliveries			
0		474	30%
1 or 2		377	24%
3 or 4		292	19%
5 or 6		207	13%
7+		212	13%
Unknown		12	1%
MDOC treated			
Prolonged/obstructive labour		848	54%
Antepartum haemorrhage		195	12%
Severe pre-eclampsia/eclampsia		165	11%
Abortum complicatum		117	7%
Post-partum haemorrhage		157	10%
Ectopic pregnancy		24	2%
Rupture uterus		30	2%
Sepsis		38	2%
Total		1574	100%

Table 3 Reasons for paediatric RS, period 2015-2017*		
Reason for referral	Number	%
Malaria	1540	30%
Anemia	910	18%
Pneumonia/ARI**	830	16%
Diarrhoea and vomiting	495	10%
Malnutrition	274	5%
Convulsion	186	4%
Hernia/Hydrocele	165	3%
Sepsis/Septicemia	127	2%
Dehydration	48	1%
Burn	30	1%
Others	522	10%
Total	5127	100%

\* For a number of patients, more than one suspected diagnosis for referral was reported; \*\* Acute Respiratory Infection.

## DISCUSSION

This study presents for the first time trends in utilization of MCH services before, during, and after Ebola, at hospital and community level from the country most heavily affected by the Ebola epidemic. It also presents data on the restructured and reorganised RS, which started immediately after the EVD outbreak. The study shows that there was a decrease in all MCH indicators and service uptake immediately after the onset of the outbreak, with a levelling or increase during the EVD period. In the post-Ebola period, all indicators (except for maternal deaths) showed an increase, in comparison with the pre-Ebola period. This was particularly marked at hospital level because the post Ebola reinforcement of the RS led to an increase in pediatric admissions, maternal admissions, and consequently a rise of institutional deliveries, C-sections, and MDOCs. In addition, while at the hospital level trends in the post-Ebola period are in line with the pre-Ebola, at community level there is a negative trend compared to the pre-Ebola period for all indicators taken into consideration. The study presents results in contrast to other studies that showed a decline in MCH services in the Ebola and post-Ebola periods.<sup>6 31 32</sup>

### Pre Ebola and Ebola periods

As mentioned above, the approach implemented in the Pujehun district<sup>17 28</sup> avoided vertical interventions only focused on the containment of the EVD epidemic. It worked on strengthening all the components of the health system - before, during, and long after the epidemic. This approach may have contributed to reducing the spread of infection and the impact of the disease on MCH services.<sup>17 18</sup> As shown by this paper, at community level family planning, ANC, and institutional deliveries, were affected only at the beginning of the Ebola outbreak with a small decrease in service utilization. In contrast, Jones et al., evaluated the number of antenatal and postnatal visits, institutional births, emergency obstetric care (EmOC), maternal deaths and stillbirths across 13 districts of Sierra Leone for 10 months during, and 12 months prior to the epidemic. They found that following the onset of the epidemic there was an 18% decrease in the number of women attending ANC visits and an 11% decrease in the number of women attending for birth at healthcare facilities.<sup>14</sup>

During the Ebola epidemic, the Pujehun hospital maintained C-sections and delivery volume at pre-Ebola levels. There was a stable number of patients attending the hospital during the Ebola outbreak, as shown by the number of maternal and pediatric admissions. The study of Brolin and colleagues focused on in-hospital deliveries and C-section volume in Sierra Leone. They showed that nationwide, albeit with substantial variation between districts, in-hospital deliveries and C-sections

1  
2  
3 364 decreased by over 20% during the Ebola outbreak, mainly because of the closure of not-for-profit  
4  
5 365 hospitals.<sup>6</sup> Brolin also noted that in general, at hospital level, in Sierra Leone those facilities that  
6  
7 366 remained open performed about the same number of deliveries and C-sections after the onset of the  
8  
9 367 EVD outbreak as they did before.<sup>6</sup> This seems to indicate that the decrease observed at national level  
10  
11 368 was related to the closing of key health facilities. The number of Ebola cases was not uniform  
12  
13 369 throughout districts in Sierra Leone and Pujehun was one of the least affected districts. The low  
14  
15 370 number of cases may also have helped to maintain public confidence in service provision and uptake  
16  
17 371 of services.<sup>7 8</sup>

17  
18 372  
19  
20 373 **Post Ebola period**

21  
22 374 There is a shortage of data in Sierra Leone and the other West Africa countries affected regarding the  
23  
24 375 resumption of services after the epidemic. Pujehun district showed contrasting results at community  
25  
26 376 level. Results of the post Ebola vs pre-Ebola show an increase of activities for institutional delivery  
27  
28 377 and ANC 4. However, there is a negative trend among the two periods, for the variables taken into  
29  
30 378 consideration, namely institutional deliveries (-7, 95% CI -10 to -4,  $p < 0.001$ ) ANC 1 (-6, 95% CI -10 to  
31  
32 379 -3,  $p < 0.001$ ), ANC 4 (-8, 95% CI -11 to -5,  $p < 0.001$ ) and family planning (-85, 95% CI -119 to -51,  $p$   
33  
34 380  $< 0.001$ ). In 2017, the Pujehun district showed a coverage of 98% for ANC 1 (98% in 2013), 91% for  
35  
36 381 ANC 4 (76% at national level in 2013) and 90% for institutional deliveries (62% in 2013).<sup>33 34</sup> The initial  
37  
38 382 intervention carried out by DWA in the period 2012-2014 at the community level probably increased  
39  
40 383 these percentages, with an initial growth of the trend that had been slowing down in the years 2016-  
41  
42 384 2017. Possible explanations for this may include: bypassing, i.e. using alternative health care instead  
43  
44 385 of free or subsidized public clinics; increased opportunities to get transport to seek healthcare in  
45  
46 386 neighbouring districts; reduced demand for MCH services at community level; and reduced quality of  
47  
48 387 MCH services at PHUs.

47 388 A study by Camara et al. in a rural district of Guinea showed a considerable recovery gap in the  
48  
49 389 post-Ebola period for ANC (37%) and institutional deliveries (34%).<sup>31</sup> Also Delamou et al. noted a  
50  
51 390 significant reduction in the average number of ANC visits and institutional deliveries during the Ebola  
52  
53 391 outbreak, in 6 districts of Guinea, and the overall post-outbreak trends did not suggest recovery.<sup>32</sup> By  
54  
55 392 contrast, Wagenaar et al., which analysed 10 primary care indicators in Liberia, before, during, and  
56  
57 393 after the Ebola outbreak, showed significant positive trends during the post-EVD period for ANC and  
58  
59 394 institutional deliveries.<sup>35</sup>

59  
60



There are multifactorial and complex reasons for the decline of family planning in the Pujehun district. The activities that MoHS and Dwa implemented from 2012 onwards were maintained during and after the EVD epidemic. However, a general decrease in the availability of healthcare personnel and international aid was observed and this could be a factor in the family planning decline. A possible stock-out of family planning methods has also been suggested as a reason for the decrease.<sup>25</sup> In addition, a reduction in demand for family planning in the post Ebola period could account for the decline of the service. Experiencing a disaster can trigger the desire to “rebuild” communities, reducing the need for family planning methods,<sup>36</sup> or communities may prefer traditional methods of contraception.<sup>37</sup> However, the reduction in family planning use in Pujehun district did not translate into an increase in institutional deliveries as occurred in neighbouring Liberia.<sup>38</sup> Although no further transmissions of Ebola took place in the Pujehun district after November 2015, the awareness of the ongoing transmission elsewhere in Sierra Leone, in Guinea and Liberia might have influenced health seeking behaviours.<sup>39 40</sup> However, this does not seem to have influenced other types of MCH services at community level. For comparison, the above mentioned study of Camara et al. showed that the utilization of family planning declined by 51% during the Ebola outbreak but recovered in the post-Ebola period.<sup>31</sup>

At hospital level, the situation is different. In the post-Ebola period, there was a significant increase in the volumes of activities: pediatric and maternal admissions, MDOC cases, deliveries, and C-sections. This increase can be directly linked to the reorganization and strengthening of the RS immediately after the Ebola epidemic. Based on the 3 delays theory,<sup>41</sup> in Pujehun it was decided to tackle the second delay, a lack of accessibility to health services. The distance to the hospital as well as lack of accessible and affordable vehicles were recognized as significant barriers when attempting to access CEmONC services at the hospital.<sup>42 43</sup> The success of the RS service can be linked to the integration of the key components needed for a successful service, namely: i) a transport system which took account of the specific geographical characteristics of the district;<sup>42</sup> ii) an effective communication system with a call center in contact with all PHUs of the district, the ambulance drivers, and the hospital; iii) training of all the PHU staff on the recognition of obstetric emergencies and on the RS.<sup>44 45</sup> Several meetings were planned with local community leaders and religious leaders to raise awareness of the importance of giving birth in health facilities. Prohibitive costs have been shown to be a major factor in preventing women accessing health facilities during childbirth in Sierra Leone.<sup>42 46 47</sup> Meetings were also organised to inform the population that the service was free of charge, and to give reassurance that the ambulances carried no risk of Ebola infection to people using



1  
2  
3 427 them. The increase in complicated cases treated at the hospital did not translate into an increase in  
4  
5 428 maternal and pediatric deaths, reflecting positively on the quality of care provided. The maternity  
6  
7 429 ward death rate remained around 1% throughout the 2012-2017 study period. The differences in  
8  
9 430 average death rates during the period 2015-2017 among referred and not referred pediatric patients  
10  
11 431 were 10.5% and 4.3% respectively. This showed that the pediatric RS works for the most critical cases  
12  
13 432 able to reach the hospital in time.

14 433  
15  
16 434 **CONCLUSIONS**

17  
18 435 There are a number of contextual factors and limitations that should be taken into account in the  
19  
20 436 analysis of the results of this study. The data refers to a single area of Sierra Leone and therefore our  
21  
22 437 sample cannot be considered representative of the country as a whole. We defined our distinct period  
23  
24 438 of EVD outbreak arbitrarily, from one month before the first case in the district to three months after  
25  
26 439 the last case in the district. This was done because the EVD crisis affected areas of the country outside  
27  
28 440 Pujehun prior to and after outbreak within Pujehun. The official end of the EVD epidemic for Sierra  
29  
30 441 Leone was declared on March 17, 2016, and for the countries of Guinea and Liberia was declared on  
31  
32 442 June 1, 2016. All the results should be taken with some degree of statistical caution, because no  
33  
34 443 correction was performed to take into account the multiplicity of the tests carried out. Finally, our  
35  
36 444 study assumed that no other interventions in addition to those described occurred concurrently with  
37  
38 445 the Ebola epidemic.<sup>18</sup> Similarly, we assumed that no other substantial interventions in addition to the  
39  
40 446 re-organisation of the RS happened in the post-Ebola period which would have affected the service  
41  
42 447 trends that we observed. The Pujehun district had 49 confirmed EVD cases. This number is much  
43  
44 448 lower than in other districts. If it is true that the fear of Ebola may have prevented people from  
45  
46 449 accessing health services, the small number of EVD cases in the community may have also raised  
47  
48 450 confidence, leading to the increase of utilization rates after the initial drop. The strength of this study  
49  
50 451 is that it uses data from a remote rural district in Sierra Leone, with a 6-year observational period.  
51  
52 452 The pre, intra, and post-Ebola periods data, allowed a comparison between trends. DwA was working  
53  
54 453 in this community before the outbreak began, which gave an advantage of knowledge of the setting  
55  
56 454 when the epidemic began, which in turn facilitated mitigating measures to be put in place. In addition,  
57  
58 455 this allowed a collection of data in a prospective way, reducing the potential bias in the accuracy of  
59  
60 456 the data reported by other studies.<sup>6 14 32 35</sup>

58 457 Failures in providing effective health care are associated with a chiefly vertical focus on outbreak  
59  
60 458 control.<sup>19-21</sup> The approach implemented in the Pujehun district worked on strengthening all the

components of the health system - governance, human resources, community involvement - before, during and, after the epidemic.

The strengthening of the health system in the district, compared to other districts, allowed the containment of the epidemic and, above all, to maintain and strengthen MCH services as shown by the data reported in the paper. Health facilities in the district, both at community and hospital level, were able to maintain their services during the epidemic, overcoming public fear of Ebola and lack of confidence in service providers, which led to the public staying away from facilities in other districts in Sierra Leone.<sup>14</sup> In post-crisis situations, "windows of opportunity" are opened for redirecting the policies of the national health systems, renovating specific sectors (e.g. human resources, epidemiological surveillance systems, financing, etc.) and renewing services/practices at the operational level.<sup>48</sup> In Pujehun the implementation of an RS immediately after the acute Ebola phase might have reduced delays in patients accessing care and enabled a significant improvement in all MCH indicators at hospital level. Other studies have also found that using this window of opportunity to introduce systems such as performance based financing can also produce positive outcomes.<sup>49</sup> As Sierra Leone continues its recovery, there is a need to quantify the impact of the outbreak on MCH care to guide long-term strategies for MHC services. This study provides evidence on strategies to increase the resilience of fragile healthcare services and the importance of NGOs and government collaboration to bring about change.

**Figure 1** Study area, the Pujehun district in Sierra Leone.

**Figure 2** Maternal and pediatric admissions at hospital level.

**Figure 3** C-sections, deliveries, MDOCs, pediatric and maternal deaths at hospital level.

**Figure 4** ANC 1, ANC 4, deliveries, and family planning at community level.

1  
2  
3 489  
4  
5 490  
6  
7 491  
8  
9 492  
10  
11 493  
12 494  
13  
14 495  
15  
16 496  
17  
18 497  
19  
20 498  
21  
22 499  
23 500  
24  
25 501  
26  
27 502  
28  
29 503  
30  
31 504  
32  
33 505  
34 506  
35  
36 507  
37  
38 508  
39  
40 509  
41  
42 510  
43  
44 511  
45 512  
46  
47 513  
48  
49 514  
50  
51 515  
52  
53 516  
54 517  
55  
56 518  
57  
58 519  
59  
60 520  
521

**Acknowledgements** We are deeply grateful to all the staff of Pujehun district hospital and Peripheral Health Units, District Health Management, and the personnel of Doctors with Africa CUAMM who worked in Pujehun during the epidemic. We thank James Dean for his help with proof-reading.

**Contributors** GLQ, FT, and GP contributed to study design, literature review, data analysis, data interpretation, writing, and review of the final manuscript. LF, FDG, DP, and CM performed data analysis, data interpretation, and drafting of the manuscript. DB, SS, AK, BZ, and VP contributed to data collection and data interpretation. BSC, AS, WM, and SJ contributed to data interpretation and review of the final manuscript. All co-authors contributed to the improvement of the article.

**Funding** This work was undertaken and funded by Doctors with Africa CUAMM.

**Competing Interests** None declared.

**Patient consent for publication** Obtained.

**Ethics approval** Sierra Leone Ethics and Scientific Review Committee, Directorate of Policy, Planning and Information, Ministry of Health and Sanitation, Sierra Leone.

**Data sharing statement** All data underlying the findings described in the manuscript are fully available without restriction. No additional data are available.

**Disclaimer** The views expressed in this publication are the sole responsibility of the authors and do not necessarily reflect the views of the affiliated organisations.

## REFERENCES

1. World Health Organization. *Ebola response roadmap. Situation reports*. WHO, 2018. Available from: <http://www.who.int/csr/disease/ebola/situation-reports/archive/en/> [Accessed March 2019].
2. Jones S, Ameh C. Exploring the impact of the Ebola outbreak on routine maternal health services in Sierra Leone, 2015. Available from: [https://www.vsointernational.org/sites/vso\\_international/files/vso\\_sierra\\_leone\\_\\_](https://www.vsointernational.org/sites/vso_international/files/vso_sierra_leone__) [Accessed March 2019].
3. Streifel C. *How did Ebola impact maternal and child health in Liberia and Sierra Leone? A report of the CSIS Global Health Policy Center*. CSIS, 2015.
4. Brolin Ribacke KJ, Saulnier DD, Eriksson A, *et al*. Effects of the West Africa Ebola Virus Disease on Health-Care Utilization - A Systematic Review. *Front Public Health* 2016;4:222.
5. Government of Sierra Leone, Ministry of Health and Sanitation. *Ebola virus disease situation report*. MoHS, 2015.
6. Brolin Ribacke KJ, van Duinen AJ, Nordenstedt H, *et al*. The Impact of the West Africa Ebola Outbreak on Obstetric Health Care in Sierra Leone. *PLoS One* 2016;11:e0150080.
7. Cancedda C, Davis SM, Dierberg KL, *et al*. Strengthening Health Systems While Responding to a Health Crisis: Lessons Learned by a Nongovernmental Organization During the Ebola Virus Disease Epidemic in Sierra Leone. *J Infect Dis* 2016; 214 (Suppl. 3):S153-S163.
8. Jones S, Sam B, Bull F, Pieh SB, *et al*. 'Even when you are afraid, you stay': Provision of maternity care during the Ebola virus epidemic: A qualitative study. *Midwifery* 2017;52:19-26.
9. Dynes MM, Miller L, Sam T, *et al*. Perceptions of the risk for Ebola and health facility use among health workers and pregnant and lactating women Kenema District, Sierra Leone, 2014. *Morb Mortal Wkly Rep* 2015;63:12267.
10. Delamou A, Hammonds RM, Caluwaerts S, *et al*. Ebola in Africa: beyond epidemics, reproductive health in crisis. *Lancet* 2014; 13; 384:2105.
11. Hayden EC. Ebola obstructs malaria control. *Nature* 2014;514:15-6.
12. Sierra Leone, Ministry of Health and Sanitation, United Nations Children's Fund. *Health facility survey. Assessing the impact of the EVD outbreak on health systems in Sierra Leone*. MoHS, UNICEF, 2014.
13. Sierra Leone, Ministry of Health and Sanitation, UN Children's Fund. *Sierra Leone health facility assessment 2015: impact of the EVD [Ebola Virus Disease] outbreak on Sierra Leone's primary health care system*. MoHS, UNICEF, 2015.

1  
2  
3 569 14. Jones S, Gopalakrishnan S, Ameh CA, *et al.* 'Women and babies are dying but not of Ebola': the  
4 570 effect of the Ebola virus epidemic on the availability, uptake and outcomes of maternal and  
5 571 newborn health services in Sierra Leone. *BMJ Glob Health* 2016;1(3):e000065.  
7 572  
8 573 15. Centers for Disease Control and Prevention (CDC). 2014-2016 Ebola Outbreak Distribution in West  
9 574 Africa. 2017. Available from:[https://www.cdc.gov/vhf/ebola/history/2014-2016-](https://www.cdc.gov/vhf/ebola/history/2014-2016-outbreak/distribution-map.html)  
10 575 [outbreak/distribution-map.html](https://www.cdc.gov/vhf/ebola/history/2014-2016-outbreak/distribution-map.html)[Accessed March 2019].  
12 576  
13 577 16. Sierra Leone, Ministry of Health and Sanitation. *Nationwide needs assessment for emergency*  
14 578 *obstetric and new-born care services in Sierra Leone*. MoHS, 2008.  
15 579  
16 580 17. Ajelli M, Parlamento S, Bome D, *et al.* The 2014 Ebola virus disease outbreak in Pujehun, Sierra  
18 581 Leone: epidemiology and impact of interventions. *BMC Med* 2015;13:281.  
19 582  
20 583 18. Quaglio GL, Pizzol D, Bome D, *et al.* Maintaining maternal and child health services during the Ebola  
21 584 outbreak: experience from Pujehun, Sierra Leone. *PLoS Currents* 2016;8:  
22 585 [ecurrents.outbreaks.d67aea257f572201f835772d7f188ba5](https://doi.org/10.1371/journal.ploscurrents.outbreaks.d67aea257f572201f835772d7f188ba5).  
24 586  
25 587 19. Dubois M, Wake C, Sturridge S, Bennett C. *The Ebola response in West Africa. Exposing the politics*  
26 588 *and culture of international aid*. 2015. Available from:  
27 589 <https://www.odi.org/sites/odi.org.uk/files/odi-assets/publications-opinion-files/9903.pdf>  
29 590 [Accessed March 2019]  
30 591  
31 592 20. Scott V, Crawford-Browne S, Sanders D. Critiquing the response to the Ebola epidemic through a  
32 593 primary health care approach. *BMC Public Health* 2016; 16:410 DOI 10.1186/s12889-016-3071-4.  
34 594  
35 595 21. Kruk ME, Myers M, Varpilah ST, *et al.* What is a resilient health system? Lessons from Ebola. *Lancet*  
36 596 2015;385:1910-12.  
37 597  
38 598 22. The Guardian. *Sierra Leone declares first Ebola free district*. 10 January 2015. Available  
39 599 from:<http://www.theguardian.com/world/2015/jan/10/sierraleonefirstebolafreedistrictwho>  
41 600 [Accessed March 2019].  
42 601  
43 602 23. Theuring S, Koroma AP, Harms G. "In the hospital, there will be nobody to pamper me": a  
44 603 qualitative assessment on barriers to facility-based delivery in post-Ebola Sierra Leone. *Reprod*  
45 604 *Health* 2018;15:155: 10.1186/s12978-018-0601-9.  
47 605  
48 606 24. Ly J, Sathananthan V, Griffiths T, *et al.* Facility-based delivery during the Ebola virus disease  
49 607 epidemic in rural Liberia: analysis from a cross-sectional, population-based household survey. *PLoS*  
50 608 *Med* 2016;13: e1002096.  
52 609  
53 610 25. Barden-O'Fallon J, Barry MA, Brodish P, *et al.* Rapid assessment of Ebola-related implications for  
54 611 reproductive, maternal, newborn, and child health service delivery and utilization in Guinea. *PLoS*  
55 612 *Curr* 2015;7:ecurrents.outbreaks.0b0ba06009d d091bc39ddb3c6d7b0826.  
57 613  
58 614 26. Lori JR, Rominski SD, Perosky JE, *et al.* A case series study on the effect of Ebola on facility-based  
59 615 deliveries in rural Liberia. *BMC Pregnancy Childbirth* 2015;15:254.  
60


27. Iyengar P, Kerber K, Howe CJ, *et al.* Services for mothers and newborns during the Ebola outbreak in Liberia: the need for improvement in emergencies. *PLoS Curr* 2015; 7:ecurrents.outbreaks.4ba318308719ac86fbef91f8e56cb66f.
28. World Health Organization. *Situation report: Ebola virus disease, 10 June 2016*. WHO, 2018. Available from: [http://apps.who.int/iris/bitstream/10665/208883/1/ebolasitrep\\_10Jun2016\\_eng.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/208883/1/ebolasitrep_10Jun2016_eng.pdf?ua=1) [Accessed March 2019]
29. World Health Organisation. Monitoring emergency obstetric care. 2009 Available from: [https://apps.who.int/iris/bitstream/handle/10665/44121/9789241547734\\_eng.pdf;jsessionid=81643CFAAD44ECAEE6445C3BCBA40026?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/44121/9789241547734_eng.pdf;jsessionid=81643CFAAD44ECAEE6445C3BCBA40026?sequence=1) [Accessed March 2019]
30. R Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. Vienna, Austria, 2018.
31. Camara BS, Delamou A, Diro E, *et al.* Effect of the 2014/2015 Ebola outbreak on reproductive health services in a rural district of Guinea: an ecological study. *Trans R Soc Trop Med Hyg* 2017;11:22-9.
32. Delamou A, Hammonds RM, Caluwaerts S, *et al.* Effect of Ebola virus disease on maternal and child health services in Guinea: a retrospective observational cohort study. *Lancet Glob Health* 2017;5:e448-e457.
33. Statistics Sierra Leone, United Nations Children's Fund. *Sierra Leone multiple indicator cluster survey 2017*. Survey findings report. Stats SL, UNICEF, 2018.
34. Sierra Leone, Ministry of Health and Sanitation. *Sierra Leone demographic and health survey*. MoHS, 2013.
35. Wagenaar BH, Augusto O, Beste J, *et al.* The 2014-2015 Ebola virus disease outbreak and primary healthcare delivery in Liberia: Time-series analyses for 2010-2016. *PLoS Med* 2018;15:e1002508.
36. Nobles J, Frankenberg E, Thomas D. The Effects of mortality on fertility: population dynamics after a natural disaster. *Demography* 2015;52:15-38.
37. Hapsari ED, Widyawati, Nisman WA, *et al.* Change in contraceptive methods following the Yogyakarta earthquake and its association with the prevalence of unplanned pregnancy. *Contraception* 2009;79:316-22.
38. McBain RK, Wickett E, Mugunga JC, *et al.* The post-Ebola baby boom: time to strengthen health systems *Lancet* 2016;388: 2331-3.
39. Bolkan HA, Bash-Taqi DA, Samai M, *et al.* Ebola and indirect effects on health service function in Sierra Leone. *PLoS Curr* 2014;6:ecurrents.outbreaks.0307d588d f619f9c9447f8ead5b72b2d.



1  
2  
3 662 40. Olu O, Kargbo B, Kamara S, *et al.* Epidemiology of Ebola virus disease transmission among health  
4 663 care workers in Sierra Leone, May to December 2014: a retrospective descriptive study. *BMC Infect*  
5 664 *Dis* 2015;15:416.  
6 665  
7 666 41. Thaddeus S, Maine D. Too far to walk: maternal mortality in context. *Sm Sci Med* 1994;38:1091-  
8 667 1110. 41.  
9 668  
10 669 42. Treacy L. Distance, accessibility and costs. Decision-making during childbirth in rural Sierra Leone:  
11 670 A qualitative study. *PLoS One* 2018;13: e0188280.  
12 671  
13 672 43. Tayler-Smith K, Zachariah R, Manzi M, *et al.* An ambulance referral network improves access to  
14 673 emergency obstetric and neonatal care in a district of rural Burundi with high maternal mortality.  
15 674 *Trop Med Int Health* 2013;18:993-1001.  
16 675  
17 676 44. Groppi L, Somigliana E, Pisani V, *et al.* A hospital-centred approach to improve emergency obstetric  
18 677 care in South Sudan. *Int J Gynaecol Obstet* 2015;128:58–61.  
19 678  
20 679 45. Tsegaye A, Somigliana E, Alemayehu T, *et al.* Ambulance referral for emergency obstetric care in  
21 680 remote settings. *Int J Gynaecol Obstet* 2016;133:316-9.  
22 681  
23 682 46. Oyerinde K, Harding Y, Amara P, *et al.* Barriers to uptake of emergency obstetric and newborn care  
24 683 services in Sierra Leone: a qualitative study. *Comm Med & Health Educ* 2012;2: 10.4172/2161-  
25 684 0711.1000149.  
26 685  
27 686 47. Vallières F, Cassidy EL, McAuliffe E, *et al.* Can Sierra Leone maintain the equitable delivery of their  
28 687 Free Health Care Initiative? The case for more contextualised interventions: results of a cross-  
29 688 sectional survey. *BMC Health Serv Res* 2016;16:258.  
30 689  
31 690 48. Witter S, Hunter B. *Resilience of health systems during and after crises – what does it mean and*  
32 691 *how can it be enhanced?* ReBUILD Research Programme Consortium. 2017. Available  
33 692 from:[https://rebuildconsortium.com/media/1535/rebuild\\_briefing\\_1\\_june\\_17\\_resilience.pdf](https://rebuildconsortium.com/media/1535/rebuild_briefing_1_june_17_resilience.pdf)  
34 693 [Accessed March 2019].  
35 694  
36 695 49. Mussah VG, Mapleh L, Ade S, *et al.* Performance-based financing contributes to the resilience of  
37 696 health services affected by the Liberian Ebola outbreak. *Public Health Action* 2017;7(Suppl.  
38 697 1):S100-S105.  
39 698  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60









1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

BMJ Open

- C-sections
- Deliveries
- Maternal deaths
- MDOCs
- Pediatric deaths





## ANNEX 1

### Statistical analysis

For each indicator, a segmented seasonal autoregressive model of order 1 was estimated. The segments defined the three periods: before the EVD epidemic (January 2012 to May 2014), during the epidemic (June 2014 to February 2015), and after the epidemic (March 2015 to December 2017). The model for each indicator  $Y_t$  collected at hospital or community level was as follows:  $Y_t = \beta_0 + \beta_1 T_t + \beta_2 X_t + \beta_3 X_t T_t + \beta_4 Z_t + \beta_5 Z_t T_t + \beta_6 \text{Month} + \varepsilon_t$ .  $\beta_0$  estimates the number of individuals using the service at the beginning of the pre-Ebola period;  $\beta_1$  estimates the average monthly change in the number using the service over the pre-outbreak period;  $T_t$  is the time since the start of the study;  $\beta_2$  represents the change in the level of service use that occurred in the period immediately after the EVD period (designated by indicator variable  $X_t$ );  $\beta_3$  represents the difference between the trend in service use during the EVD outbreak compared to the pre-disease period;  $\beta_4$  represents the change in service use that occurred in the period immediately after the end of the outbreak (post-outbreak period designated by indicator variable  $Z_t$ );  $\beta_5$  is the difference between the trend in service use during the period after the Ebola virus disease outbreak compared with the period during the outbreak period;  $\beta_m$  represents a series of indicator variables for each calendar month, and  $t$  is the random error term. Overall trends across the periods and the comparisons among trends were calculated as follows: linear trend during the outbreak =  $\beta_1 + \beta_3$ ; linear trend after the outbreak =  $\beta_1 + \beta_3 + \beta_5$ ; and linear trend after the outbreak vs linear trend before the outbreak =  $\beta_3 + \beta_5$ . Average levels across the periods and their comparisons were calculated as follows: average during the outbreak =  $\beta_0 + \beta_2$ ; average after the outbreak =  $\beta_0 + \beta_2 + \beta_4$ ; and difference between after the outbreak and before the outbreak =  $\beta_2 + \beta_4$ . Differences were considered statistically significant at  $p < 0.05$ . The analysis was performed using R.<sup>30</sup>

ANNEX 2.

Full results analysis.

Maternal admissions, maternal deaths, C-sections, and MDOCs at hospital level.

	Maternal admissions			Maternal deaths			C-sections			MDOC		
	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value
Difference between average of Ebola period vs pre-Ebola period	7	-7 to 22	0.333	-1	-2 to 0	0.042	5	-1 to 11	0.13	2	-11 to 14	0.782
Difference between average of post-Ebola period vs Ebola period	43	28 to 58	<0.001	2	1 to 3	0.001	15	8 to 21	<0.001	41	30 to 54	<0.001
Difference between average of post-Ebola period vs pre-Ebola period	50	37 to 64	<0.001	1	0 to 2	0.135	19	13 to 25	<0.001	43	31 to 54	<0.001
Pre-Ebola period												
Number of events over pre-Ebola period ( $\beta_0$ )	49	37 to 61	<0.001	1	0 to 2	0.026	9	4 to 14	0.001	16	5 to 26	0.003
Trend in number over pre-Ebola period ( $\beta_1$ )	0	0 to 1	0.281	0	0 to 0	0.677	0	-0 to 0	0.999	0	0 to 0.5	0.768
Ebola period												
Average monthly change in number over Ebola period ( $\beta_2$ )	-40	-60 to -19	<0.001	0	-2 to 0	0.480	2	-7 to 11	0.668	-11	-29 to 6	0.207
Difference between trend of Ebola period vs pre-Ebola period ( $\beta_3$ )	7	4 to 11	<0.001	0	0 to 0	0.605	1	-1 to 2	0.346	4	1 to 7	0.006
Post-Ebola period												
Average monthly change in number during post-Ebola period ( $\beta_4$ )	11	-7 to 30	0.23	1	0 to 2	0.258	13	5 to 21	0.001	16	0 to 32	0.044
Difference between trend of post-Ebola period vs Ebola period ( $\beta_5$ )	-7	-10 to -4	<0.001	0	0 to 0	0.665	-1	-2 to 0.8	0.433	-4	-7 to -1	0.009
Difference between trend of post-Ebola vs pre-Ebola period ( $\beta_3 + \beta_5$ )	0	-1 to 1	1	0	0 to 0	0.657	0	0 to 0	0.431	0	0 to 1	0.503

Pediatric admissions, pediatric deaths, and institutional deliveries at hospital level.

	Pediatric admissions			Pediatric deaths			Institutional deliveries		
	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value
Difference between average of Ebola period vs pre-Ebola period	1	-39 to 40	0.968	-1	-6 to 5	0.826	11	2 to 21	0.02
Difference between average of post-Ebola period vs Ebola period	133	92 to 174	<0.001	9	3 to 15	0.004	28	18 to 38	<0.001
Difference between average of post-Ebola period vs pre-Ebola period	134	98 - 170	<0.001	8	3 to 14	0.003	39	31 to 48	<0.001
Pre-Ebola period									
Number of events over pre-Ebola period ( $\beta_0$ )	46	10 to 82	0.011	7	2 to 12	0.007	27	19 to 34	<0.001
Trend in number over pre-Ebola period ( $\beta_1$ )	0	-2 to 2	0.808	0	0 to 0	0.641	0	0 to 0	0.42
Ebola period									
Average monthly change in number over Ebola period ( $\beta_2$ )	1	-48 to 50	0.955	1	-7 to 9	0.836	-12	-25 to 1	0.072
Difference between trend of Ebola period vs pre-Ebola period ( $\beta_3$ )	1	-8 to 10	0.823	0	-1 to 2	0.763	4	2 to 6	0.001
Post Ebola period									
Average monthly change in number over post-Ebola period ( $\beta_4$ )	53	5 to 100	0.029	6	-1 to 14	0.086	11	-1 to 22	0.064
Difference between trend of post-Ebola period vs Ebola period ( $\beta_5$ )	2	-7 to 10	0.702	0	-1 to 1	0.899	-3	-5 to -1	0.001
Difference between trend of post-Ebola vs pre-Ebola period ( $\beta_3 + \beta_5$ )	3	0 to 5	0.035	0	0 to 0	0.423	0	0 to 0	0.486

Institutional delivery, ANC 1, ANC 4 and family planning at community level.

	Institutional delivery			ANC 1			ANC 4			Family planning		
	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value	$\beta$	95% CI	p value
Difference between average of Ebola period vs pre-Ebola period	148	99 to 196	<0.001	74	3 to 145	0.042	80	21 to 139	0.008	490	-92 to 1073	0.099
Difference between average of post-Ebola period vs Ebola period	-10	-59 to 39	0.695	-48	-122 to 26	0.2	23	-38 to 84	0.461	-262	-855 to 330	0.386
Difference between average of post-Ebola period vs pre-Ebola period	138	93 to 183	<0.001	26	-40 to 91	0.448	103	48 to 157	<0.001	228	-293 to 750	0.391
Pre Ebola period												
Number of events over pre-Ebola period ( $\beta_0$ )	688	643 to 732	<0.001	1062	1002 to 1121	<0.001	694	644 to 743	<0.001	2690	2187 to 3193	<0.001
Trend in number over pre-Ebola period ( $\beta_1$ )	8	6 to 10	<0.001	7	4 to 10	<0.001	6	4 to 8	<0.001	69	42 to 95	<0.001
Ebola period												
Average monthly change in number over Ebola period ( $\beta_2$ )	-28	-90 to 34	0.382	-61	-161 to 40	0.238	-94	-176 to -11	0.027	-671	-1431 to 89	0.084
Difference between trend of Ebola period vs pre-Ebola period ( $\beta_3$ )	-1	-12 to 10	0.881	-5	-21 to 12	0.591	5	-8 to 19	0.437	-26	-156 to 104	0.692
Post Ebola period												
Average monthly change in number during post-Ebola period ( $\beta_4$ )	-25	-81 to 30	0.37	-5	-94 to 83	0.906	35	-37 to 109	0.343	-51	-759 to 657	0.888
Difference between trend of post-Ebola period vs Ebola period ( $\beta_5$ )	-7	-17 to 4	0.228	-2	-18 to 15	0.819	-13	-27 to 0	0.056	-59	-186 to 68	0.361
Difference between trend of post-Ebola vs pre-Ebola period ( $\beta_3 + \beta_5$ )	-7	-10 to -4	<0.001	-6	-10 to -3	<0.001	-8	-11 to -5	<0.001	-85	-119 to -51	<0.001

## STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No.	Recommendation	Page No.	Relevant text from manuscript
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1	
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2	
<b>Introduction</b>				
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4-6	
Objectives	3	State specific objectives, including any prespecified hypotheses	6	
<b>Methods</b>				
Study design	4	Present key elements of study design early in the paper	7	
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	7-8	
Participants	6	(a) <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	8	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6	
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	7-8	
Bias	9	Describe any efforts to address potential sources of bias	16-17	
Study size	10	Explain how the study size was arrived at		

Continued on next page

Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	9-10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	10
		(b) Describe any methods used to examine subgroups and interactions	10
		(c) Explain how missing data were addressed	NA
		(d) <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy NA	NA
		(e) Describe any sensitivity analyses	NA
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	9-12
		(b) Give reasons for non-participation at each stage	NA
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	7
		(b) Indicate number of participants with missing data for each variable of interest	
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	9-11
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	9-11
		(b) Report category boundaries when continuous variables were categorized	9-11
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	9-11

Continued on next page

Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	NA
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	13
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	16-17
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	16-17
Generalisability	21	Discuss the generalisability (external validity) of the study results	16-17
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	18

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).